

Rec'd PCT/PTO 14 APR 2005

R E N C H R E P U B L I C

INPI

NATIONAL INSTITUTE
FOR INDUSTRIAL
PROPERTY

PATENT OF INVENTION

UTILITY CERTIFICATE - CERTIFICATE OF ADDITION

OFFICIAL COPY

The Director General of the National Institute for Industrial Property certifies that the attached document is the true certified copy of an application for an Industrial Property Right filed at the Institute.

Issued in Paris, 21 JUL. 2003

For the Director General of the
National Institute for Industrial Property,
The Head of the Patents Department

(signature)

Martine PLANCHE

NATIONAL
INSTITUTE FOR
INDUSTRIAL
PROPERTY

NATIONAL PUBLIC INSTITUTION

DB 267/141102

HEAD OFFICE

26 bis, rue de Saint Petersburg
75800 PARIS cedex 08
Telephone: 33 (0)1 53 04 53 04
Facsimile: 33 (0)1 53 04 45 23
www.inpi.fr

CREATED BY LAW NO. 51-444 OF 19th APRIL 1951

INPI

**National
Institute for
Industrial Property**
26bis, rue de Saint Pétersbourg
75800 Paris Cedex 08
Telephone: 01 53 04 53 04
Facsimile: 01 42 94 86 54

**PATENT OF INVENTION
UTILITY CERTIFICATE**
Intellectual Property Code - Book VI

cerfa
No. 11354*01

REQUEST FOR GRANT 1/2

Reserved for INPI

This form is to be completed legibly in black ink DB 540 W /260899

DEPOSITION OF DOCUMENTS DATE 18 OCT. 2002 PLACE 75 INPI PARIS NATIONAL REGISTRATION NO. GIVEN BY THE INPI 0212965 FILING DATE GIVEN BY THE INPI 18 OCT. 2002		1 NAME AND ADDRESS OF THE APPLICANT OR OF THE AUTHORISED AGENT TO WHOM CORRESPONDENCE MUST BE ADDRESSED LES LABORATOIRES SERVIER Direction Brevets 12, Place de La Défense 92415 COURBEVOIE Cedex FRANCE	
Your references for this file (optional) 29681			
Confirmation of a deposit by facsimile		<input type="checkbox"/> No. given by INPI to the facsimile	
2 NATURE OF THE APPLICATION		Mark one of the following 4 boxes	
Patent application		<input checked="" type="checkbox"/>	
Application for a Utility Certificate		<input type="checkbox"/>	
Divisional application <i>Initial patent application or initial utility certificate application</i>		<input type="checkbox"/> No. _____ Date _____ No. _____ Date _____	
Conversion of a European Patent Application <i>Initial patent application</i>		<input type="checkbox"/> No. _____ Date _____	
3 TITLE OF THE INVENTION (maximum 200 characters or spaces) New substituted [1,4]benzodioxino[2,3-e]isoindole compounds, a process for their preparation and pharmaceutical compositions containing them.			
4 DECLARATION OF PRIORITY OR REQUEST FOR THE BENEFIT OF THE FILING DATE OF A PRIOR FRENCH APPLICATION		Country or organisation _____ Date _____ No. _____ Country or organisation _____ Date _____ No. _____ Country or organisation _____ Date _____ No. _____ <input type="checkbox"/> If there are other priorities, mark the box and use the "Continuation" form	
5 APPLICANT		<input type="checkbox"/> If there are other Applicants, mark the box and use the "Continuation" form	
Surname or company name		LES LABORATOIRES SERVIER	
Forenames			
Legal nature			
SIREN No.			
APE-NAF Code			
Address	Street	12, Place de La Défense	
	Postal code and town	92415	COURBEVOIE Cedex
Country		FRANCE	
Nationality		FRENCH	
Telephone no. (optional)		01.55.72.60.00	
Facsimile no. (optional)		01.55.72.72.13	
E-mail address (optional)			

INPINational
Institute for
Industrial Property

First filing

**PATENT OF INVENTION
UTILITY CERTIFICATE**

REQUEST FOR GRANT 2/2

Reserved for INPI

DEPOSITION OF DOCUMENTS	
DATE 18 OCT. 2002	
PLACE 75 INPI PARIS	
NATIONAL REGISTRATION NO. GIVEN BY THE INPI 0212965	
DB 540 W /260899	
Your references for this file: (optional) 29681	
6 AUTHORISED AGENT	
Surname WENGER	
Forename Sabine	
Practice or company LES LABORATOIRES SERVIER	
No. of standing power of attorney and/or of contractual bond	
Address	
Street 12, Place de La Défense	
Postal code and town 92415 COURBEVOIE Cedex	
Telephone no. (optional) 01.55.72.60.00	
Facsimile no. (optional) 01.55.72.72.13	
E-mail address (optional)	
7 INVENTOR(S)	
The inventors are the Applicants <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No In this case, supply a separate designation of inventorship	
8 SEARCH REPORT	
For a patent application only (including division and conversion)	
immediate drawing up <input checked="" type="checkbox"/> or deferred drawing up <input type="checkbox"/>	
Payment of the fees in instalments	
Payment in three instalments, for natural persons only <input type="checkbox"/> Yes <input type="checkbox"/> No	
9 REDUCTION IN FEES	
For natural persons only <input type="checkbox"/> Requested for the first time for this invention (attach a notice of non-imposition) <input type="checkbox"/> Requested prior to this deposit (attach a copy of the admissibility decision for this invention or indicate its reference)	
If you have used the "Continuation" form, indicate the number of pages attached	
10 SIGNATURE OF THE APPLICANT OR OF THE AUTHORISED AGENT (Name and position of signatory)	
Sabine WENGER, Patent Engineer (signature)	
STAMP OF THE PREFECTURE OR OF THE INPI	
[signature]	


Law No. 78-17 of 6 January 1978 relating to information processing, data files and rights applies to the responses made on this form. It guarantees right of access to and correction of the data concerning you at the INPI.

The present invention relates to new substituted [1,4]benzodioxino[2,3-e]isoindole compounds, to a process for their preparation and to pharmaceutical compositions containing them. Use of the compounds of the present invention is promising because of their anti-tumour activity.

5 Patent Application WO 00/18407 describes pyrrolocarbazole compounds for use in the prevention and treatment of deafness and the sensation of loss of balance. Patent Applications US 5 705 511 and WO 96/11933 describe cyclopenta[g]pyrrolo[3,4-e]indole compounds fused at their indole moiety and cyclopentene moiety to an aromatic or non-
10 aromatic cyclic system optionally containing hetero atoms. Those compounds possess pharmacological activities making them of use especially in the treatment of cancerous cells. Patent Application WO 01/85686 describes pyrrolocarbazole-aryl compounds for use in the treatment of cancers. Patent Application EP 0 841 337 claims substituted 7,12-dioxo-benzo[a]anthracene compounds and describes their anti-cancer properties.



15 The compounds of the present invention are novel in terms of both their structure and their use as anti-tumour agents.

More specifically, the present invention relates to compounds of formula (I) :



wherein :

- A, together with the carbon atoms to which it is bonded, represents a group of formula (a) or (b) :



wherein :

- ✧ W_1 , together with the carbon atoms to which it is bonded, represents a phenyl group or a pyridyl group,
- 5 ✧ Z represents a group selected from hydrogen and halogen atoms and the groups linear or branched (C_1-C_6) alkyl, nitro, cyano, hydroxy, linear or branched (C_1-C_6) alkoxy, aryl, aryl- (C_1-C_6) alkyl (in which the alkyl moiety is linear or branched), aryloxy and aryl- (C_1-C_6) alkoxy (in which the alkoxy moiety is linear or branched) and NR_5R_6 wherein R_5 and R_6 , which are identical or different, each

10 independently of the other represents a group selected from a hydrogen atom and a linear or branched (C_1-C_6) alkyl group,
- ✧ R_4 represents a group selected from a hydrogen atom and the groups linear or branched (C_1-C_6) alkyl, aryl and aryl- (C_1-C_6) alkyl (in which the alkyl moiety is linear or branched) or a group $-C(O)-OR'_5$ wherein R'_5 represents a group selected from the

15 groups linear or branched (C_1-C_6) alkyl, aryl and aryl- (C_1-C_6) alkyl (in which the alkyl moiety is linear or branched),
- Y represents a group selected from an oxygen atom and a methylene group,
- R_2 represents a hydrogen atom and, in that case:

R₃ represents a group selected from a hydrogen atom and the groups linear or branched (C₁-C₆)alkyl, aryl, aryl-(C₁-C₆)alkyl (in which the alkyl moiety is linear or branched) and SO₂CF₃,

- or R₂ and R₃ form a bond,

- 5
- R₁ represents a group selected from a hydrogen atom and the groups linear or branched (C₁-C₆)alkyl, aryl and aryl-(C₁-C₆)alkyl (in which the alkyl moiety is linear or branched) or a linear or branched (C₁-C₆)alkylene chain substituted by one or more identical or different groups selected from -OR"₅ and -NR"₅R"₆ wherein R"₅ and R"₆ are as defined for R₅ and R₆ defined hereinbefore,

- 10
- Z₁ and Z₂, which are identical or different, each independently of the other represent a hydrogen atom or
Z₁ and Z₂, together with the carbon atoms carrying them, form a phenyl group,

with the proviso that, when Z represents a hydrogen atom, R₁ is other than a hydrogen atom,

- 15
- to their enantiomers, diastereoisomers, N-oxide, and to addition salts thereof with a pharmaceutically acceptable acid or base,


- 20
- wherein "aryl" is to be understood as meaning a phenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, indenyl or indanyl group, each of those groups being optionally substituted by one or more identical or different groups selected from halogen, linear or branched (C₁-C₆)alkyl, linear or branched (C₁-C₆)trihaloalkyl, hydroxy, linear or branched (C₁-C₆)alkoxy, and amino optionally substituted by one or two linear or branched (C₁-C₆)alkyl groups.

- 25
- Among the pharmaceutically acceptable acids there may be mentioned, without implying any limitation, hydrochloric acid, hydrobromic acid, sulphuric acid, phosphonic acid, acetic acid, trifluoroacetic acid, lactic acid, pyruvic acid, malonic acid, succinic acid,

glutaric acid, fumaric acid, tartaric acid, maleic acid, citric acid, ascorbic acid, methane-sulphonic acid, camphoric acid etc..


Among the pharmaceutically acceptable bases there may be mentioned, without implying any limitation, sodium hydroxide, potassium hydroxide, triethylamine etc..

- 5 Preferred compounds of the invention are compounds of formula (I) that correspond more especially to formula (IA) :




wherein R_1 , R_4 , W_1 , Z , Z_1 and Z_2 are as defined for formula (I).

- 10 According to a second advantageous embodiment, preferred compounds of the invention are compounds of formula (I) that correspond more especially to formula (IB) :




wherein R_1 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I).

According to a third advantageous embodiment, preferred compounds of the invention are compounds of formula (I) corresponding more especially to formula (IC) :




5 wherein R_1 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I).

According to a fourth advantageous embodiment, preferred compounds of the invention are compounds of formula (I) that correspond more especially to formula (ID) :




wherein R_1 , R_3 , R_4 , W_1 , Z , Z_1 and Z_2 are as defined for formula (I).

According to a fifth advantageous embodiment, preferred compounds of the invention are compounds of formula (I) that correspond more especially to formula (IE) :




wherein R_1 , R_3 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I).

5 According to a sixth advantageous embodiment, preferred compounds of the invention are compounds of formula (I) that correspond more especially to formula (IF) :




wherein R_1 , R_3 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I).

10 According to a seventh advantageous embodiment, preferred compounds of the invention are compounds of formula (I) that correspond more especially to formula (IG) :




wherein R_1 , R_4 , W_1 , Z , Z_1 and Z_2 are as defined for formula (I).

According to an eighth advantageous embodiment, preferred compounds of the invention are compounds of formula (I) that correspond more especially to formula (IH) :



wherein R_1 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I).


According to a ninth advantageous embodiment, preferred compounds of the invention are compounds of formula (I) that correspond more especially to formula (IJ) :



wherein R_1 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I).

In one embodiment of interest, the group Z to which preference is given in accordance with the invention is a hydrogen atom, halogen atom or hydroxy group.

Advantageously, the group A , together with the carbon atoms to which it is bonded, to which preference is given in accordance with the invention, is a group of formula :



5

In an especially advantageous embodiment, the group R_3 to which preference is given in accordance with the invention is a hydrogen atom or a linear or branched (C_1-C_6) alkyl group.

10 In an especially advantageous embodiment, the group R_1 to which preference is given in accordance with the invention is a hydrogen atom, a linear or branched (C_1-C_6) alkyl group or a linear or branched (C_1-C_6) alkylene chain substituted by one or more identical or different groups selected from $-NR_5R_6$ wherein R_5 and R_6 are as defined for formula (I).


In another embodiment of interest, the groups Z_1 and Z_2 to which preference is given in accordance with the invention are hydrogen atoms.

Compounds to which preference is given in accordance with the invention are :

- 7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 5 • 10-fluoro-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 11-fluoro-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 7-[2-(dimethylamino)ethyl]-10-fluoro[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-carbazole-6,8-dione,
- 10-hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 10 • 11-hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 7-[2-(dimethylamino)ethyl][1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 7-[2-(dimethylamino)ethyl]-10-hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-carbazole-6,8-dione,
- 7-[2-(dimethylamino)ethyl]-11-hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-carbazole-6,8-dione,
- 15 • 7-[2-(dimethylamino)ethyl][1,4]benzodioxino[2,3-e]pyrido[2',3':5,6][1,4]oxazino-[3,2-g]isoindole-6,8-dione.


The enantiomers, diastereoisomers, N-oxides, and addition salts with a pharmaceutically acceptable acid or base, of the preferred compounds form an integral part of the invention.

20 The invention relates also to a process for the preparation of compounds of formula (I), which process is characterised in that there is used as starting material a compound of formula (II) :




wherein Z_1 and Z_2 are as defined for formula (I),

which compound of formula (II) is reacted with N-bromosuccinimide in the presence of benzoyl peroxide to yield the compound of formula (III) :




wherein Z_1 and Z_2 are as defined hereinbefore,

5 which compound of formula (III) is reacted with sodium iodide to yield the compound of formula (IV) :



wherein Z_1 and Z_2 are as defined hereinbefore,


10 which compound of formula (IV) is reacted with n-butyllithium and then with trimethyltin chloride to yield the compound of formula (V) :




wherein Z_1 and Z_2 are as defined hereinbefore,

which compound of formula (V) is :


- either treated, in the presence of tetrakis(triphenylphosphine)palladium(0), with a compound of formula (VI) :




wherein P_G represents an amine-protecting group well-known in organic synthesis and W_1 and Z are as defined for formula (I), to yield the compound of formula (VII) :



- 5 wherein P_G , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
which compound of formula (VII) is treated with a compound of formula (VIII) :




wherein R_{1a} represents a hydrogen atom or a methyl group, to yield the compound of formula (IX) :




wherein P_G , R_{1a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compound of formula (IX) is treated with N-bromosuccinimide and benzoyl peroxide to yield the compound of formula (I/a), a particular case of the compounds of formula (I) :




wherein P_G , R_{1a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,


which compound of formula (I/a) is optionally treated with a compound of formula (X) :



wherein R_{1b} has the same definition as R_1 in formula (I) but is other than a hydrogen atom or a methyl group, to yield the compound of formula (I/b), a particular case of the compounds of formula (I) :




- 5 wherein R_{1b} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
which compounds of formulae (I/a) and (I/b) constitute the compounds of formula (I/c) :




wherein R'_4 represents a hydrogen atom or a group P_G , and R_1 , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

- or treated in the presence of bis(triphenylphosphine)palladium(II) chloride with a compound of formula (XI) :




wherein Boc represents a tert-butoxycarbonyl group and R_{1a}, W₁ and Z are as defined
 5 hereinbefore, to yield the compound of formula (XII) :




wherein Boc, R_{1a}, Z, Z₁, Z₂ and W₁ are as defined hereinbefore,
 which compound of formula (XII) is :

- ◆ either irradiated with a UV lamp, in the presence of iodine, in a non-polar and aprotic solvent, to yield the compounds of formulae (I/d) and (I/e), particular cases of the compounds of formula (I) :




(I/d)



(I/e),

wherein Boc, R_{1a}, Z, Z₁, Z₂ and W₁ are as defined hereinbefore,
which compounds of formula (I/d) :


- 5 ✧ optionally are subjected to deprotection of the amine function according to conventional methods of organic synthesis to yield the compound of formula (I/f), a particular case of the compounds of formula (I) :




(I/f),

wherein R_{1a}, Z, Z₁, Z₂ and W₁ are as defined hereinbefore,

- 10 ✧ or optionally are subjected to the action of a compound of formula (XIII) :




wherein R_{3a} has the same definition as R_3 in formula (I) but is other than a hydrogen atom and G is as defined hereinbefore, to yield the compound of formula (I/g), a particular case of the compounds of formula (I) :




5 wherein Boc, R_{1a} , R_{3a} , Z, Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compounds of formulae (I/d), (I/e) and (I/g) constitute the compound of formula (I/h) :




wherein Boc, R_{1a} , R_3 , Z, Z_1 , Z_2 and W_1 are as defined hereinbefore,

10 which compound of formula (I/h) optionally is subjected to the same reaction conditions as the compound of formula (I/a) to yield the compound of formula (I/i), a particular case of the compounds of formula (I) :



wherein R_{1b} , R_3 , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

♦ or subjected to the action of hydrochloric acid to yield the compound of formula (XIV) :



5


wherein R_{1a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compound of formula (XIV) is subjected to the action of a compound of formula (XV) :





10

wherein G represents a leaving group and R_{4a} has the same definition as R_4 in formula (I) but is other than a hydrogen atom, to yield the compound of formula (XVI) :




wherein R_{1a} , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compound of formula (XVI) is subjected to the same reaction conditions as the compound of formula (XII) to yield the compounds of formulae (I/j) and (I/k), particular cases of the compounds of formula (I) :




wherein R_{1a} , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compound of formula (I/k) optionally is subjected to the action of a compound of formula (XIII) as defined hereinbefore to yield the compound of formula (I/l) :




wherein R_{1a} , R_{3a} , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
which compounds of formulae (I/j), (I/k) and (I/l) constitute the compounds of
formula (I/m) :




5

wherein R_{1a} , R_3 , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
which compound of formula (I/m) optionally is subjected to the same reaction conditions
as the compound of formula (I/h) to yield the compound of formula (I/n) :



wherein R_{1b} , R_3 , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,


which compounds of formulae (I/e), (I/h) and (I/i), (I/m) and (I/n) constitute the compounds of formula (I/o) :



5 wherein R_1 , R_3 , R_4 , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,


which compounds of formulae (I/a) to (I/o) constitute the totality of the compounds of formula (I), which are purified, where necessary, according to conventional purification techniques, which may be separated, if desired, into their different isomers according to a conventional separation technique, and which are converted, if desired, into their addition salts with a pharmaceutically acceptable acid or base.

The compounds of formula (VI) may advantageously be obtained starting from a compound of formula (A) :



wherein Z and W_1 are as defined hereinbefore,


the amine function of which compound of formula (A) is protected by a protecting group P_G well-known to the person skilled in the art, to yield the compound of formula (B) :



wherein P_G represents a tert-butoxycarbonyl or phenoxycarbonyl group and W_1 and Z are as defined hereinbefore,


which compound of formula (B) is treated with lithium diisopropylamide followed by diphenyl chlorophosphate to yield the compound of formula (VI).

The compounds of formula (XI) may advantageously be obtained starting from a compound of formula (C) :




wherein R_{1a} is as defined hereinbefore,

which compound of formula (C) is treated, in the presence of lithium bis(trimethylsilyl)amide, with a compound of formula (D) :



wherein W_1 and Z are as defined hereinbefore, to yield a compound of formula (E) :



wherein R_{1a} , Z and W_1 are as defined hereinbefore,

which compound of formula (E) is reacted with *di*-tert-butyl dicarbonate in the presence of 4-dimethylaminopyridine to yield the compound of formula (XI).

- 5 The compounds of formulae (II), (VIII), (X), (XIV), (XVI), (A), (C) and (D) either are commercial compounds or are obtained according to conventional methods of organic synthesis readily accessible to the person skilled in the art.

The compounds of formula (I) have valuable pharmacological properties. They have excellent cytotoxicity *in vitro* not only with respect to leukaemia lines but also with respect
10 to solid-tumour lines; they also exert an action on the cell cycle and are active *in vivo*, in a leukaemia model. Those properties enable them to be used therapeutically as anti-tumour agents.

The present invention relates also to pharmaceutical compositions comprising compounds of formula (I), their optical isomers or an addition salt thereof with a pharmaceutically
15 acceptable base or acid, alone or in combination with one or more inert, non-toxic excipients or carriers.

Among the pharmaceutical compositions according to the invention there may be mentioned more especially those that are suitable for oral, parenteral, nasal, rectal, perlingual, ocular or respiratory administration, and especially tablets or dragées,
20 sublingual tablets, sachets, paquets, gelatin capsules, glossettes, lozenges, suppositories,

creams, ointments, dermal gels, injectable or drinkable preparations, aerosols, eye drops and nose drops.

By virtue of the pharmacological properties that are characteristic of the compounds of formula (I), the pharmaceutical compositions comprising the said compounds of formula (I) as active ingredient are, accordingly, especially useful in the treatment of cancers.

The useful dose varies according to the age and weight of the patient, the administration route, the nature of the therapeutic indication and any associated treatments and ranges from 0.1 to 400 mg per day, in one or more administrations.

The Examples that follow illustrate the invention, without limiting it in any way. The starting materials used are known products or are prepared according to known procedures.

The structures of the compounds described in the Examples were determined according to customary spectrophotometric techniques (infra-red, nuclear magnetic resonance, mass spectrometry, ...).

PREPARATION A : Benzo[1,4]dioxin-2-yl(trimethyl)stannane

Step A : 2,3-Dibromo-2,3-dihydro-1,4-benzodioxine

Under an inert atmosphere, 73.45 mmol of 1,4-benzodioxane, 150 ml of carbon tetrachloride and then 163.32 mmol of recrystallised N-bromosuccinimide and 88.1 μ mol of benzoyl peroxide are mixed together. The reaction mixture is refluxed with the aid of a lamp (60 W) for 6 hours. The precipitated succinimide is removed by filtration and the filtrate is concentrated, allowing the expected product to be isolated.

Step B : Benzo[1,4]dioxine

73.45 mmol of the compound obtained in Step A above are dissolved in 125 ml of acetone; the solution is then stirred for 2 hours under reflux, in the presence of sodium iodide (359.9 mmol). After evaporating off the solvent, the residue is dissolved in a mixture of water/ethyl acetate (100 ml/200 ml). The organic phase is then washed with 20 % aqueous sodium thiosulphate solution and then the aqueous phase is extracted twice with ethyl acetate. The organic phases are combined, dried over magnesium sulphate and then concentrated to dryness. Purification by chromatography over silica gel (petroleum ether/ethyl acetate : 95/5) allows the expected product to be obtained.

IR (NaCl film) : $\nu_{(\text{C}=\text{C enol ether})} = 1665 \text{ cm}^{-1}$; $\nu_{(\text{C}=\text{C Ar})} = 1590 \text{ cm}^{-1}$.

Step C : Benzo[1,4]dioxin-2-yl(trimethyl)stannane

14.91 mmol of 1,4-benzodioxine are dissolved in 18 ml of anhydrous tetrahydrofuran under an anhydrous atmosphere, and the temperature of the solution is lowered to -78°C . 23.86 mmol of a 1.5M solution of n-butyllithium in hexane are added dropwise to the reaction mixture, which is stirred at -78°C for 2 hours 15 minutes. 37.28 mmol of trimethyltin chloride dissolved in 10 ml of anhydrous tetrahydrofuran are added dropwise and the reaction mixture is stirred at -78°C for 45 minutes and then returned to ambient temperature for 15 hours. The reaction mixture is hydrolysed with 15 % aqueous potassium fluoride solution and is stirred for 45 minutes. The precipitated tin salts are removed by filtration, and the aqueous phase is extracted 3 times with ethyl acetate. The organic phases are combined and concentrated. Purification by flash chromatography over silica gel (petroleum ether/ethyl acetate : 95/5) allows the expected product to be obtained.

IR (NaCl film) : $\nu_{(\text{C}=\text{C enol ether})} = 1639 \text{ cm}^{-1}$; $\nu_{(\text{C}=\text{C Ar})} = 1592 \text{ cm}^{-1}$.

PREPARATION B : Naphtho[2,3-b][1,4]dioxin-2-yl(tributyl)stannane**Step A : Ethyl 2,3-dihydronaphtho[2,3-b][1,4]dioxine-2-carboxylate**

Under an inert atmosphere, 47.03 mmol of dry potassium carbonate are added to a solution of 62.43 mmol of 2,3-dihydroxynaphthalene in 150 ml of anhydrous acetone. The reaction mixture is heated at reflux, after adding 17.2 mmol of ethyl 2,3-dibromopropanoate. 15 minutes later, 47.03 mmol of dry potassium carbonate, and also 17.2 mmol of ethyl 2,3-dibromopropanoate are again added to the reaction mixture. This operation is repeated two more times, at 15-minute intervals. Refluxing is maintained for 18 hours. The reaction mixture is then filtered, and the residue is washed with acetone. The filtrate is concentrated and is then taken up in ethyl acetate and washed with 100 ml of water. The aqueous phase is extracted three times with ethyl acetate, and then the organic phases are combined, dried over magnesium sulphate, filtered and evaporated under reduced pressure. Purification by flash chromatography over silica gel (petroleum ether/ethyl acetate : 9/1) allows the expected product to be isolated.

IR (NaCl film) : $\nu_{\text{C=O}} = 1759 \text{ cm}^{-1}$.

Mass spectrum : 259 $[\text{M} + 1]^+$.

Step B : Ethyl naphtho[2,3-b][1,4]dioxine-2-carboxylate

Under an inert atmosphere, a solution of 25.94 mmol of the compound obtained in Step A above in 120 ml of carbon tetrachloride is heated at reflux with the aid of a lamp (60 W) in the presence of 57.03 mmol of recrystallised N-bromosuccinimide and a catalytic amount of benzoyl peroxide. Stirring is maintained at reflux for 2 hours 30 minutes. After cooling, the liberated succinimide is filtered off and the filtrate is concentrated to dryness *in vacuo*. The dibrominated ester obtained is dissolved in 100 ml of acetone, and then 129.62 mmol of sodium iodide are added. Stirring is maintained at ambient temperature for 4 hours. After evaporating off the solvent, the residue is taken up in a mixture of water/ethyl acetate and washed with 1M aqueous sodium thiosulphate solution. The organic phase is dried over magnesium sulphate, filtered and evaporated. Purification by flash chromatography

over silica gel (petroleum ether/ethyl acetate : 9/1) allows the expected product to be isolated.

Melting point : 98°C.

IR (KBr) : $\nu_{\text{C=O}} = 1724 \text{ cm}^{-1}$; $\nu_{\text{C=C}} = 1682 \text{ cm}^{-1}$.

5 Mass spectrum : 257 $[\text{M} + 1]^+$.

Step C : Naphtho[2,3-b][1,4]dioxine-2-carboxylic acid

10 A solution of 24.58 mmol of the compound obtained in Step B above, in 25 ml of methanol, is heated at reflux for one hour, in the presence of 20 ml of 1M aqueous lithium hydroxide solution. After cooling, and evaporating off the methanol, the reaction mixture is acidified with 1M aqueous hydrochloric acid solution until a pH = 1 is obtained. The precipitate formed is filtered off, allowing the expected product to be isolated.

Melting point : > 360°C.

IR (KBr) : $\nu_{\text{C=O}} = 1678 \text{ cm}^{-1}$; $\nu_{\text{C=C}} = 1661 \text{ cm}^{-1}$; $\nu_{\text{OH}} = 3450 \text{ cm}^{-1}$.

Mass spectrum : 229 $[\text{M} + 1]^+$.

15 **Step D : Naphtho[2,3-b][1,4]dioxine**

20 A solution of 0.66 mmol of the compound obtained in Step C above, in 1 ml of quinoline, is heated at 220°C for 3 hours, in the presence of a catalytic amount of powdered copper. After cooling, the residue is taken up in ethyl acetate and washed with 1M aqueous hydrochloric acid solution. The organic phase is dried over magnesium sulphate, filtered and concentrated. Purification by flash chromatography over silica gel (petroleum ether/ethyl acetate : 9/1) allows the expected product to be obtained.

Melting point : 96-98°C.

IR (KBr) : $\nu_{\text{C=C}} = 1665 \text{ cm}^{-1}$; $\nu_{\text{C=O}} = 1296 \text{ cm}^{-1}$.

Step E : Naphtho[2,3-b][1,4]dioxin-2-yl(tributyl)stannane

The expected product is obtained according to the procedure described in Step C of Preparation A, starting from the compound of Step D above and tributyltin.

IR (NaCl film) : $\nu_{C-O} = 1166, 1247 \text{ cm}^{-1}$.

5 Mass spectrum : 473 $[M + 1]^+$.

PREPARATION C : Phenyl 3-[(diphenoxyphosphoryl)oxy]-4H-1,4-benzoxazine-4-carboxylate**Step A : Phenyl 2,3-dihydro-4H-1,4-benzoxazin-3-one-4-carboxylate**

10 Under an anhydrous atmosphere, a solution of 10 mmol of 2H-1,4-benzoxazin-3-one in 50 ml of tetrahydrofuran is cooled to -78°C . At that temperature, 11 mmol of a 1.6M solution of *n*-butyllithium in hexane are added dropwise. After being in contact for 30 minutes at -78°C , 11 mmol of phenyl chloroformate are added dropwise and stirring is carried out for a further 2 hours. After returning to ambient temperature, the solution is hydrolysed and then extracted with ethyl acetate. The organic phase is dried over
15 magnesium sulphate, filtered and evaporated. After purification by chromatography over silica gel (petroleum ether/ethyl acetate : 8/2), the expected product is isolated.

IR (NaCl film) : $\nu_{C=O} = 1739, 1796 \text{ cm}^{-1}$.

Mass spectrum : 270 $(M + 1)$.

Step B : Phenyl 3-[(diphenoxyphosphoryl)oxy]-4H-1,4-benzoxazine-4-carboxylate

20 Under an anhydrous atmosphere, a solution of 10 mmol of the product obtained in Step A above, in 50 ml of anhydrous THF, is cooled to -78°C . At that temperature, 12 mmol of 2M LDA (in a solution of heptane/THF) are added dropwise. After stirring for 2 hours, 12 mmol of diphenyl chlorophosphate are added dropwise to the reaction mixture, which is kept for a further 2 hours at -78°C . After returning to ambient temperature, the solution is
25 hydrolysed and then extracted with ethyl acetate. The organic phase is dried over

magnesium sulphate, filtered and concentrated. After purification of the residue by chromatography over silica gel (petroleum ether/ethyl acetate : 9/1), the expected product is isolated.

IR (NaCl film) : $\nu_{\text{C=O}} = 1748 \text{ cm}^{-1}$; $\nu_{\text{P=O}} = 1315 \text{ cm}^{-1}$.

5 Mass spectrum : 502 (M + 1).

PREPARATION D : Phenyl 3-[(diphenoxyphosphoryl)oxy]-2,3-dihydro-4H-pyrido-[3,2-b][1,4]oxazine-4-carboxylate

Step A : Phenyl 2,3-dihydro-4H-pyrido[3,2-b][1,4]oxazine-4-carboxylate

10 The expected product is obtained according to the procedure described in Step A of Preparation C, using phenyl chloroformate as substrate.

Melting point : 97°C.

IR (KBr) : $\nu_{\text{C=O}} = 1717 \text{ cm}^{-1}$; 1803 cm^{-1} .

Mass spectrum : m/z 271 (M + 1).

15 **Step B : Phenyl 3-[(diphenoxyphosphoryl)oxy]-2,3-dihydro-4H-pyrido[3,2-b][1,4]-oxazine-4-carboxylate**

The expected product is obtained according to the procedure described in Step B of Preparation C, starting from the compound of the Step above.

Melting point : 82°C.

IR (KBr) : $\nu_{\text{C=O}} = 1749 \text{ cm}^{-1}$; $\nu_{\text{P=O}} 1294 \text{ cm}^{-1}$.

20 Mass spectrum : m/z 503 (M + 1).

PREPARATION E : 3-Bromo-4-(1H-1-carboxylate of tert-butyl-indol-3-yl)-1-methyl-1H-pyrrole-2,5-dione

Step A : 3-Bromo-4-(1H-indol-3-yl)-1-methyl-1H-pyrrole-2,5-dione

11.16 mmol of indole are dissolved in 29 ml of anhydrous tetrahydrofuran under an inert atmosphere. The temperature of the reaction mixture is lowered to -15°C , and then 17.85 mmol of a 1M solution of bis(trimethylsilyl)lithium in hexane are added dropwise. After stirring at -15°C for 1 hour 10 minutes, 7.44 mmol of 2,3-dibromo-*N*-methylmaleimide dissolved in 8 ml of anhydrous tetrahydrofuran are added to the reaction mixture, which is then stirred for 20 minutes at -15°C , and for 15 minutes from -15°C to ambient temperature. The reaction mixture is hydrolysed with a few millilitres of aqueous 0.3N hydrochloric acid solution until a pH of about 7 is obtained. A few millilitres of ethyl acetate are then added, and the aqueous phase is extracted four times with ethyl acetate. The organic phases are combined, washed with saturated aqueous sodium chloride solution and then evaporated. The residue is washed, filtered and rinsed with methanol, allowing the expected product to be isolated.

Melting point : 167°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}} = 1715, 1767 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3304 \text{ cm}^{-1}$.

Mass spectrum : 305 $[\text{M} + 1]^+$.

Step B : 3-Bromo-4-(1H-1-carboxylate of tert-butyl-indol-3-yl)-1-methyl-1H-pyrrole-2,5-dione

8.19 mmol of the compound obtained in Step A above are dissolved, at 0°C , in 37.5 ml of anhydrous tetrahydrofuran, in the presence of 28.68 mmol of di-*tert*-butyl dicarbonate and 614.5 μmol of 4-dimethylaminopyridine. The reaction mixture is brought to ambient temperature and stirred for 2 hours 30 minutes. After evaporating off the solvent, the residue obtained is taken up in methanol, filtered, rinsed with methanol and then dried *in vacuo*, allowing the expected product to be isolated.

Melting point : 142°C .

IR (KBr) : $\nu_{\text{C=O}} = 1762 \text{ cm}^{-1}$.

Mass spectrum : 405 $[\text{M} + 1]^+$.

PREPARATION F : 3-Bromo-4-(1*H*-indol-3-yl)-1*H*-pyrrole-2,5-dione

15.69 mmol of indole are dissolved in 25 ml of anhydrous tetrahydrofuran under an inert atmosphere. 15.69 mmol of a 3M solution of ethylmagnesium bromide in diethyl ether are added, and the reaction mixture is heated at 60°C for 1 hour 30 minutes. Once it has cooled, the mixture is added dropwise to a solution of 3.92 mmol of 2,3-dibromomaleimide in 6 ml of anhydrous tetrahydrofuran, and then the reaction mixture is stirred at ambient temperature for one hour. The reaction mixture is hydrolysed using 20 ml of 0.3N aqueous hydrochloric acid solution until a pH of 8.5 is obtained, and a few millilitres of ethyl acetate are added. The aqueous phase is extracted three times with ethyl acetate; the organic phases are then combined, washed with saturated aqueous sodium chloride solution and then evaporated to dryness *in vacuo*. The residue is washed, filtered and rinsed with methanol, allowing the expected product to be isolated.

Melting point : > 300°C.

IR (KBr) : $\nu_{\text{C=O}} = 1772 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3343, 3699 \text{ cm}^{-1}$.

Mass spectrum : 291 $[\text{M} + 1]^+$.

PREPARATION G : 3-Bromo-4-(5-fluoro-1*H*-indol-3-yl)-1-methyl-1*H*-pyrrole-2,5-dione

The expected product is obtained according to the procedure described in Step A of Preparation E, using 5-fluoroindole.

Melting point : > 300°C.

IR (KBr) : $\nu_{\text{C=O}} = 1705 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3358 \text{ cm}^{-1}$.

Mass spectrum : 323 $[\text{M} + 1]^+$.

PREPARATION H : 3-Bromo-4-(6-fluoro-1*H*-indol-3-yl)-1-methyl-1*H*-pyrrole-2,5-dione

The expected product is obtained according to the procedure described in Step A of Preparation E, using 6-fluoroindole.

5 Melting point : 201°C.

IR (KBr) : $\nu_{\text{C=O}} = 1701, 1767 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3312 \text{ cm}^{-1}$.

Mass spectrum : 323 [M + 1]⁺.

PREPARATION I : 3-Bromo-4-(5-fluoro-1*H*-indol-3-yl)-1*H*-pyrrole-2,5-dione

10 The expected product is obtained according to the procedure described in Preparation F, using 5-fluoroindole.

Melting point : 219°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}} = 1723, 1769 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3358 \text{ cm}^{-1}$.

Mass spectrum : 309,5 [M + 1]⁺.

PREPARATION J : 3-Bromo-4-(6-fluoro-1*H*-indol-3-yl)-1*H*-pyrrole-2,5-dione

15 The expected product is obtained according to the procedure described in Preparation F, using 6-fluoroindole.

Melting point : 199°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}} = 1723, 1778 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3329 \text{ cm}^{-1}$.

Mass spectrum : 309 [M + 1]⁺.

PREPARATION K : *tert*-Butyl 5-(benzyloxy)-3-(4-bromo-1-methyl-2,5-dioxo-2,5-dihydro-1*H*-pyrrol-3-yl)-1*H*-indole-1-carboxylate

Step A : 3-[5-(Benzyloxy)-1*H*-indol-3-yl]-4-bromo-1-methyl-1*H*-pyrrole-2,5-dione

The expected product is obtained according to the procedure described in Step A of Preparation E, using 5-benzyloxyindole.

Melting point : 150°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}} = 1698 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3315 \text{ cm}^{-1}$.

Mass spectrum : 411 [M + 1]⁺.

Step B : *tert*-Butyl 5-(benzyloxy)-3-(4-bromo-1-methyl-2,5-dioxo-2,5-dihydro-1*H*-pyrrol-3-yl)-1*H*-indole-1-carboxylate

The expected product is obtained according to the procedure described in Step B of Preparation E, starting from the compound of Step A above.

Melting point : 155°C.

IR (KBr) : $\nu_{\text{C=O}} = 1709, 1738, 1773 \text{ cm}^{-1}$.

Mass spectrum : 511 [M + 1]⁺.

PREPARATION L : *tert*-Butyl 6-(benzyloxy)-3-(4-bromo-1-methyl-2,5-dioxo-2,5-dihydro-1*H*-pyrrol-3-yl)-1*H*-indole-1-carboxylate

Step A : 3-[6-(Benzyloxy)-1*H*-indol-3-yl]-4-bromo-1-methyl-1*H*-pyrrole-2,5-dione

The expected product is obtained according to the procedure described in Step A of Preparation E, using 6-benzyloxyindole.

Melting point : 138°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}} = 1705, 1762 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3314 \text{ cm}^{-1}$.

Mass spectrum : 411 [M + 1]⁺.

Step B : *tert-Butyl 6-(benzyloxy)-3-(4-bromo-1-methyl-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl)-1H-indole-1-carboxylate*

The expected product is obtained according to the procedure described in Step B of Preparation E, starting from the compound of Step A above.

5 Melting point : 158°C (decomposition).

IR (KBr) : $\nu_{C=O} = 1715, 1737, 1745 \text{ cm}^{-1}$.

Mass spectrum : 511 $[M + 1]^+$.

PREPARATION M : *tert-Butyl 5-(benzyloxy)-3-[4-bromo-1-(tert-butoxycarbonyl)-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl]-1H-indole-1-carboxylate*

10 **Step A** : *3-[5-(Benzyloxy)-1H-indol-3-yl]-4-bromo-1H-pyrrole-2,5-dione*

The expected product is obtained according to the procedure described in Preparation F, using 5-benzyloxyindole.

Melting point : 154°C (decomposition).

IR (KBr) : $\nu_{C=O} = 1697 \text{ cm}^{-1}$; $\nu_{NH} = 3333 \text{ cm}^{-1}$.

15 Mass spectrum : 397 $[M + 1]^+$.

Step B : *tert-Butyl 5-(benzyloxy)-3-[4-bromo-1-(tert-butoxycarbonyl)-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl]-1H-indole-1-carboxylate*

The expected product is obtained according to the procedure described in Step B of Preparation E, starting from the compound of Step A above.

20 Melting point : 134°C.

IR (KBr) : $\nu_{C=O} = 1743, 1768, 1801 \text{ cm}^{-1}$.

Mass spectrum : 495 $[M - \text{Boc}]^+$.

PREPARATION N : *tert*-Butyl 6-(benzyloxy)-3-[4-bromo-1-(*tert*-butoxycarbonyl)-2,5-dioxo-2,5-dihydro-1*H*-pyrrol-3-yl]-1*H*-indole-1-carboxylate

Step A : 3-[6-(Benzyloxy)-1*H*-indol-3-yl]-4-bromo-1*H*-pyrrole-2,5-dione

The expected product is obtained according to the procedure described in Preparation F,
5 using 6-benzyloxyindole.

Melting point : 166°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}}$ = 1697, 1762 cm^{-1} ; ν_{NH} = 3353 cm^{-1} .

Mass spectrum : 397 $[\text{M} + 1]^+$.

Step B : *tert*-Butyl 6-(benzyloxy)-3-[4-bromo-1-(*tert*-butoxycarbonyl)-2,5-dioxo-2,5-dihydro-1*H*-pyrrol-3-yl]-1*H*-indole-1-carboxylate
10

The expected product is obtained according to the procedure described in Step B of Preparation E, starting from the compound of Step A above.

Melting point : 120°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1719, 1744, 1764, 1807 cm^{-1} .

PREPARATION O : *tert*-Butyl 7-(benzhydryloxy)-3-(4-bromo-1-methyl-2,5-dioxo-2,5-dihydro-1*H*-pyrrol-3-yl)-1*H*-indole-1-carboxylate
15

Step A : 1-(Benzhydryloxy)-2-nitrobenzene

To a solution of dry ortho-nitrophenol (79 mmol) in 200 ml of acetone there are added 127 mmol of potassium carbonate and also 79 mmol of diphenylmethane bromide. The
20 reaction mixture is heated at reflux for 6 hours and, after the temperature has been lowered to ambient temperature, is stirred further overnight. The reaction mixture is filtered and rinsed with acetone. The filtrate is evaporated and then taken up in diethyl ether and

hydrolysed. The aqueous phase is extracted with diethyl ether; the organic phase is then dried over magnesium sulphate, filtered and concentrated. The residue obtained is taken up in petroleum ether, filtered, rinsed with petroleum ether and then dried *in vacuo*, allowing the expected product to be isolated.

5 Melting point : 98°C.

Mass spectrum : 323 $[M + NH_4^+]^+$.

Step B : 7-(Benzhydryloxy)-1H-indole

11.95 mmol of the compound obtained in Step A above are dissolved, under an inert atmosphere, in 80 ml of anhydrous tetrahydrofuran. The temperature of the reaction mixture is then lowered to -40°C and then 41.84 mmol of a 1M solution of vinylmagnesium bromide in tetrahydrofuran are added dropwise to the solution, which is then stirred for 2 hours 50 minutes from -40°C to 0°C. The reaction mixture is then hydrolysed at 0°C using 100 ml of a saturated aqueous ammonium chloride solution, and the aqueous phase is extracted three times with ethyl acetate. The organic phases are combined, dried over magnesium sulphate, filtered and evaporated. Purification by flash chromatography over silica gel (petroleum ether/ethyl acetate : 9/1) allows the expected product to be isolated.

Melting point : 114°C (decomposition).

IR (HBR) : $\nu_{NH} = 3425 \text{ cm}^{-1}$.

20 Mass spectrum : 300 $[M + 1]^+$.

Step C : 3-[7-(Benzhydryloxy)-1H-indol-3-yl]-4-bromo-1-methyl-1H-pyrrole-2,5-dione

The expected product is obtained according to the procedure described in Step A of Preparation E, using the compound of Step B above.

Melting point : 205°C.

25 IR (KBr) : $\nu_{C=O} = 1699, 1763 \text{ cm}^{-1}$; $\nu_{NH} = 3345 \text{ cm}^{-1}$.

Mass spectrum : 487 $[M + 1]^+$.

Step D : *tert-Butyl 7-(benzhydryloxy)-3-(4-bromo-1-methyl-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl)-1H-indole-1-carboxylate*

The expected product is obtained according to the procedure described in Step B of Preparation E, starting from the compound of Step C above.

Melting point : 148°C (decomposition).

IR (KBr) : $\nu_{C=O} = 1709, 1759 \text{ cm}^{-1}$.

Mass spectrum : 587 $[M + 1]^+$.

EXAMPLE 1 : *Phenyl 7-methyl-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]phenoxazine-15(6H)carboxylate*

Step A : *Phenyl 3-(1,4-benzodioxin-2-yl)-4H-1,4-benzoxazine-4-carboxylate*

Under an inert atmosphere, a mixture of 1 mmol of the compound of Preparation C, 2 mmol of the compound of Preparation A, 3 mmol of lithium chloride and tetrakis-(triphenylphosphine)palladium(0) 5 % in 5 ml of tetrahydrofuran is heated at reflux. After cooling and concentrating, the residue is taken up in ethyl acetate and washed with saturated sodium chloride solution. The organic phase is dried over magnesium sulphate, filtered and concentrated. Purification by chromatography over silica gel (petroleum ether/ethyl acetate : 6/4) allows the expected product to be isolated.

Melting point : 75°C (gum).

IR (KBr) : $\nu_{C=O} = 1739 \text{ cm}^{-1}$.

Mass spectrum : 386 $[M + 1]^+$.

Step B : *Phenyl 7-methyl-6,8-dioxo-5b,6,7,8,8a,8b-hexahydro[1,4]benzo-dioxino[2,3-a]pyrrolo[3,4-c]phenoxazine-15(5aH)-carboxylate*

In a closed system, 1 mmol of the compound obtained in Step A above and 3 mmol of *N*-methylmaleimide are stirred at 95°C for 2 hours in the presence of a few drops of toluene.

5 Purification by chromatography over silica gel (petroleum ether/ethyl acetate : 3/7) allows the expected product to be obtained.

Melting point : > 250°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1709, 1769 cm^{-1} .

Mass spectrum : 497 $[\text{M} + 1]^+$.

10 **Step C :** *Phenyl 7-methyl-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]phenoxazine-15(6H)-carboxylate*

Under an inert atmosphere, 1 mmol of the compound obtained in Step B above and 3 mmol of recrystallised *N*-bromosuccinimide are heated in distilled carbon tetrachloride at reflux for 7 minutes with the aid of a lamp (60 W), in the presence of a catalytic amount of benzoyl peroxide. After cooling, the solution is filtered and concentrated. Purification by chromatography over silica gel (petroleum ether/ethyl acetate : 1/1) allows the expected product to be obtained.

Melting point : 130°C (gum).

IR (KBr) : $\nu_{\text{C=O}}$ = 1714, 1745 cm^{-1} .

20 Mass spectrum : 493 $[\text{M} + 1]^+$.

EXAMPLE 2 : *7-[2-(Dimethylamino)ethyl][1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-phenoxazine-6,8(7H,15H)-dione*

Under an inert atmosphere, 0.1 mmol of the compound of Example 1 in 2 ml of *N,N*-dimethylethylenediamine is heated at 100°C for 6 hours. After cooling, the solution is

concentrated. Purification by chromatography over silica gel (dichloromethane/methanol : 8/2) allows the expected product to be obtained.

Melting point : 150°C.

IR (KBr) : $\nu_{C=O}$ = 1701, 1751 cm^{-1} ; ν_{NH} = 3423 cm^{-1} .

5 Mass spectrum : 430 $[M + 1]^+$.

EXAMPLE 3: Phenyl 7-methyl-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-e]-pyrido[2',3':5,6][1,4]oxazino[3,2-g]isoindole-15(6*H*)-carboxylate

Step A: Phenyl 3-(1,4-benzodioxin-2-yl)-4*H*-pyrido[3,2-b][1,4]oxazine-4-carboxylate

10 The expected product is obtained according to the procedure described in Step A of Example 1, starting from the compound of Preparation D.

Melting point : 142°C.

IR (KBr) : $\nu_{C=O}$ = 1741 cm^{-1} .

Mass spectrum : 387 $[M + 1]^+$.

15 **Step B:** Phenyl 7-methyl-6,8-dioxo-5*b*,6,7,8,8*a*,8*b*-hexahydro[1,4]benzodioxino-[2,3-*e*]pyrido[2',3':5,6][1,4]oxazino[3,2-*g*]isoindole-15(5*aH*)-carboxylate

The expected product is obtained according to the procedure described in Step B of Example 1, starting from the compound of Step A above.

Melting point : 245°C.

20 IR (KBr) : $\nu_{C=O}$ = 1709, 1769 cm^{-1} .

Mass spectrum : 498 $[M + 1]^+$.

Step C : *Phenyl 7-methyl-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-e]pyrido-[2',3':5,6][1,4]oxazino[3,2-g]isoindole-15(6H)-carboxylate*

The expected product is obtained according to the procedure described in Step C of Example 1, starting from the compound obtained in Step B above.

5 Melting point : > 250°C.

IR (KBr) : $\nu_{C=O}$ = 1711, 1760 cm^{-1} .

Mass spectrum : 494 $[M + 1]^+$.

EXAMPLE 4 : *7-[2-(Dimethylamino)ethyl][1,4]benzodioxino[2,3-e]pyrido[2',3':5,6]-[1,4]oxazino[3,2-g]isoindole-6,8(7H,15H)-dione*

10 The expected product is obtained according to the procedure of Example 2, starting from the compound of Example 3.

Melting point : 150°C (decomposition).

IR (KBr) : $\nu_{C=O}$ = 1703, 1759 cm^{-1} ; ν_{NH} = 3433 cm^{-1} .

Mass spectrum : 431 $[M + 1]^+$.

15 **EXAMPLE 5 :** *tert-Butyl 7-methyl-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-13(6H)-carboxylate*

Step A : *tert-Butyl 3-[4-(1,4-benzodioxin-2-yl)-1-methyl-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl]-1H-indole-1-carboxylate*

20 To a solution of 592.23 μmol of the compound of Preparation A in 10 ml of 1,4-dioxane there are added 493.52 μmol of the compound of Preparation E, 49.35 μmol of copper iodide and 49.35 μmol of tetrakis(triphenylphosphine)palladium. The reaction mixture is heated at 100°C for 3 hours 35 minutes, under an inert atmosphere. The solution is then filtered in order to remove the remnants of palladium, and the filtrate is evaporated.

Purification by flash chromatography over silica gel (petroleum ether/ethyl acetate/triethylamine : 8/1.9/0.1) allows the expected product to be obtained.

Melting point : 140°C.

IR (KBr) : $\nu_{C=O}$ = 1704, 1740 cm^{-1} .

5 Mass spectrum : 459 $[M + 1]^+$.

Step B : *tert-Butyl 7-methyl-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-13(6H)-carboxylate*

218.1 μmol of the compound obtained in Step A above and 5.91 mmol of double-sublimed iodine are dissolved in 500 ml of toluene inside the cell of an irradiation apparatus. After
10 irradiating for 30 minutes, with stirring, the reaction mixture is cooled and then diluted with ethyl acetate. Washing with 80 ml of 20 % aqueous sodium thiosulphate solution is carried out until the organic phase is decolourised. The aqueous phase is extracted with ethyl acetate three times, and then the organic phases are combined and directly concentrated *in vacuo*. A brown, semi-oily, semi-solid residue is collected. The procedure
15 is repeated a further five times in the same manner, using the same amounts. The crude reaction products of the six procedures are combined. Washing with ethyl acetate is carried out on the six crude reaction products, which are then filtered. The filtrate is evaporated and then taken up in methanol, filtered, rinsed with methanol and then dried *in vacuo*, allowing the expected product to be isolated.

20 Melting point : 189°C (decomposition).

Mass spectrum : 457 $[M + 1]^+$.

EXAMPLE 6 : 7-Methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-(7H,13H)-dione

Step A : 3-(1,4-Benzodioxin-2-yl)-4-(1H-indol-3-yl)-1-methyl-1H-pyrrole-2,5-dione

331.54 μmol of the compound of Step A of Example 5 are dissolved in 9 ml of formic acid. The reaction mixture is stirred under an inert atmosphere and at ambient temperature for 3 hours 30 minutes. The solvent is removed under reduced pressure; the residue is then washed, filtered and rinsed with methanol and is then dried *in vacuo*, allowing the expected product to be isolated.

Melting point : 219°C.

IR (KBr) : $\nu_{\text{C=O}} = 1697 \text{ cm}^{-1}$.

Mass spectrum : 359.5 $[\text{M} + 1]^+$.

Step B : 7-Methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione

418.5 μmol of the compound obtained in Step A above are dissolved in 50 ml of toluene in an irradiation reactor and then 5.91 mmol of double-sublimed iodine are introduced into the reaction mixture. Irradiation is carried out at 500 W for 1 hour 10 minutes. The reaction mixture is then diluted with ethyl acetate, after which it is washed with 80 ml of 20 % aqueous sodium thiosulphate solution until the organic phase is decolourised. The aqueous phase is extracted three times with ethyl acetate; the organic phases are then combined and directly concentrated. The solid residue obtained is washed firstly with diethyl ether and then filtered. It is washed subsequently with ethyl acetate and then with tetrahydrofuran and filtered. Purification by flash chromatography over silica gel (petroleum ether/ethyl acetate : 6/4) allows the expected product to be obtained.

Melting point : > 355°C.

IR (KBr) : $\nu_{\text{C=O}} = 1692, 1703 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3343 \text{ cm}^{-1}$.

Mass spectrum : 357 $[\text{M} + 1]^+$.

EXAMPLE 7 : 7-[2-(Dimethylamino)ethyl][1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-carbazole-6,8(7H,13H)-dione

238.5 μmol of the compound of Example 6 are dissolved in 4 ml of *N,N*-dimethylethylenediamine. The reaction mixture is heated at reflux for 16 hours and it is then cooled and concentrated. The residue obtained is washed, filtered and rinsed with methanol before being dried *in vacuo*, allowing the expected product to be isolated.

Melting point : 298°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}}$ = 1703, 1753 cm^{-1} ; ν_{NH} = 3441 cm^{-1} .

Mass spectrum : 414 $[\text{M} + 1]^+$.

EXAMPLE 8 : 7,13-Dimethyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-(7H,13H)-dione

Step A : 3-(1,4-Benzodioxin-2-yl)-1-methyl-4-(1-methyl-1H-indol-3-yl)-1H-pyrrole-2,5-dione

69.76 μmol of the compound obtained in Step A of Example 6, 111.62 μmol of sodium hydroxide, 7.46 μmol of benzyltriethylammonium chloride and 418.58 mmol of methyl iodide are dissolved in 0.5 ml of dichloromethane. The reaction mixture is stirred at ambient temperature for 1 hour 45 minutes, and then the solvent is removed *in vacuo*. Purification by flash chromatography over silica gel (from petroleum ether/ethyl acetate : 7/3 to ethyl acetate) allows the expected product to be obtained.

Melting point : 204°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}}$ = 1697 cm^{-1} .

Mass spectrum : 373 $[\text{M} + 1]^+$.

Step B : *7,13-Dimethyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-(7H,13H)-dione*

The expected product is obtained according to the procedure described in Step B of Example 5, starting from the compound obtained in Step A above.

5 **EXAMPLE 9 :** *13-Ethyl-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione*

Step A : *3-(1,4-Benzodioxin-2-yl)-4-(1-ethyl-1H-indol-3-yl)-1-methyl-1H-pyrrole-2,5-dione*

10 The expected product is obtained according to the procedure described in Step A of Example 8, starting from the compound obtained in Step A of Example 6 and replacing the methyl iodide with ethyl iodide.

Step B: *13-Ethyl-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8(7H,13H)dione*

15 The expected product is obtained according to the procedure described in Step B of Example 6, starting from the compound obtained in Step A above.

EXAMPLE 10 : *Naphtho[2',3':5,6][1,4]dioxino[2,3-a]pyrrolo[3,4-c]carbazole-5,7-(6H,16H)-dione*

Step A : *3-(1H-Indol-3-yl)-4-naphtho[2,3-b][1,4]dioxin-2-yl-1H-pyrrole-2,5-dione*

20 The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compounds of Preparations B and F.

Melting point : 261°C.

IR (KBr) : $\nu_{C=O} = 1759 \text{ cm}^{-1}$; $\nu_{NH} = 3171, 3398 \text{ cm}^{-1}$.

Mass spectrum : 395 [M + 1]⁺.

Step B : *Naphtho[2',3':5,6][1,4]dioxino[2,3-a]pyrrolo[3,4-c]carbazole-5,7(6H,16H)-dione*

5 The expected product is obtained according to the procedure described in Step B of Example 6, starting from the compound obtained in the Step above.

EXAMPLE 11 : *10-Fluoro-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione*

Step A : *3-(1,4-Benzodioxin-2-yl)-4-(5-fluoro-1H-indol-3-yl)-1-methyl-1H-pyrrole-2,5-dione*

10 The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation G.

Melting point : 229°C.

IR (KBr) : $\nu_{\text{C=O}} = 1698 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3380 \text{ cm}^{-1}$.

Mass spectrum : 377 [M + 1]⁺.

15 **Step B :** *10-Fluoro-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione*

The expected product is obtained according to the procedure described in Step B of Example 6, starting from the compound obtained in Step A above.

Melting point : > 355°C.

20 IR (KBr) : $\nu_{\text{C=O}} = 1699, 1753 \text{ cm}^{-1}$; $\nu_{\text{NH}} = 3339 \text{ cm}^{-1}$.

Mass spectrum : 375 [M + 1]⁺.

EXAMPLE 12 : 7-[2-(Dimethylamino)ethyl]-10-fluoro[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione

The expected product is obtained according to the procedure described in Example 7, starting from the compound of Example 11.

5 Melting point : 337°C (decomposition).

IR (KBr) : $\nu_{C=O} = 1701, 1755 \text{ cm}^{-1}$; $\nu_{NH} = 3447 \text{ cm}^{-1}$.

Mass spectrum : 432 $[M + 1]^+$.

EXAMPLE 13 : 11-Fluoro-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione

10 Step A : 3-(1,4-Benzodioxin-2-yl)-4-(6-fluoro-1H-indol-3-yl)-1-methyl-1H-pyrrole-2,5-dione

The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation H.

Melting point : 214°C.

15 IR (KBr) : $\nu_{C=O} = 1693 \text{ cm}^{-1}$; $\nu_{NH} = 3322 \text{ cm}^{-1}$.

Mass spectrum : 377 $[M + 1]^+$.

Step B : 11-Fluoro-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-(7H,13H)-dione

20 The expected product is obtained according to the procedure described in Step B of Example 6, starting from the compound obtained in Step A above.

Melting point : > 360°C.

IR (KBr) : $\nu_{C=O} = 1697, 1752 \text{ cm}^{-1}$; $\nu_{NH} = 3349 \text{ cm}^{-1}$.

Mass spectrum : 375 $[M + 1]^+$.

EXAMPLE 14 : 7-[2-(Dimethylamino)ethyl]-11-fluoro[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione

The expected product is obtained according to the procedure described in Example 7, starting from the compound of Example 13.

5 Melting point : 301°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}}$ = 1703, 1752 cm^{-1} ; ν_{NH} = 3439 cm^{-1} .

Mass spectrum : 432 $[\text{M} + 1]^+$.

EXAMPLE 15 : 10-Fluoro[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-(7H,13H)-dione

10 Step A : *3-(1,4-Benzodioxin-2-yl)-4-(5-fluoro-1H-indol-3-yl)-1H-pyrrole-2,5-dione*

The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation I.

Melting point : 241°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}}$ = 1693 cm^{-1} ; ν_{NH} = 3177, 3335 cm^{-1} .

15 Mass spectrum : 363 $[\text{M} + 1]^+$.

Step B : *10-Fluoro[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione*

The expected product is obtained according to the procedure described in Step B of Example 6, starting from the compound obtained in Step A above.

20 Melting point : > 360°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1707 cm^{-1} ; ν_{NH} = 3258, 3379 cm^{-1} .

Mass spectrum : 361 $[\text{M} + 1]^+$.

EXAMPLE 16 : 11-Fluoro[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-(7H,13H)-dione

Step A : *3-(1,4-Benzodioxin-2-yl)-4-(6-fluoro-1H-indol-3-yl)-1H-pyrrole-2,5-dione*

5 The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation J.

Melting point : 203°C.

IR (KBr) : $\nu_{C=O}$ = 1697, 1762 cm^{-1} ; ν_{NH} = 3179, 3344 cm^{-1} .

Mass spectrum : 363 $[M + 1]^+$.

10 **Step B :** *11-Fluoro[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-(7H,13H)-dione*

The expected product is obtained according to the procedure described in Step B of Example 6, starting from the compound obtained in Step A above.

Melting point : > 360°C.

IR (KBr) : $\nu_{C=O}$ = 1710 cm^{-1} ; ν_{NH} = 3190, 3455 cm^{-1} .

15 Mass spectrum : 361 $[M + 1]^+$.

EXAMPLE 17 : tert-Butyl 10-(benzyloxy)-7-methyl-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-13(6H)-carboxylate

Step A : *tert-Butyl 3-[4-(1,4-benzodioxin-2-yl)-1-methyl-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl]-5-(benzyloxy)-1H-indole-1-carboxylate*

20 The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation K.

Melting point : 93°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1703, 1706, 1737, 1741 cm^{-1} .

Mass spectrum : 565.5 $[\text{M} + 1]^+$.

5 **Step B :** **tert-Butyl 10-(benzyloxy)-7-methyl-6,8-dioxo-7,8-dihydro[1,4]-
benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-13(6H)-carboxylate**

177.1 μmol of the compound obtained in Step A above and 5.91 mmol of double-sublimed iodine are dissolved in 500 ml of toluene inside the cell of an irradiation apparatus. After irradiating for 40 minutes at 500 W, with stirring, the reaction mixture is cooled and then diluted with ethyl acetate. Washing with 80 ml of 20 % aqueous sodium thiosulphate
10 solution is carried out until the organic phase is decolourised. The aqueous phase is extracted with ethyl acetate three times, and then the organic phases are combined and directly concentrated *in vacuo*. The procedure is repeated a further two times in the same manner, using the same amounts. The crude reaction products of the three procedures are combined and the brown solid obtained is washed, filtered and rinsed with methanol before
15 being dried *in vacuo*, allowing the expected product to be isolated.

Melting point : 302°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}}$ = 1705, 1731, 1760 cm^{-1} .

Mass spectrum : 463 $[\text{M-Boc} + 1]^+$.

20 **EXAMPLE 18 :** **10-Hydroxy-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-
carbazole-6,8(7H,13H)-dione**

44.4 μmol of the compound of Example 17 are dissolved, under an inert atmosphere, in 3.5 ml of anhydrous dichloromethane. A 1M solution of boron tribromide in dichloromethane (89 μmol) is then added dropwise and with vigorous stirring to the solution, which has been cooled to 0°C. The reaction mixture is stirred at 0°C for
25 15 minutes and then for 1 hour 15 minutes whilst being allowed to return to ambient temperature. After hydrolysis with distilled water, the reaction mixture is diluted with ethyl acetate, and the aqueous phase is extracted three times with ethyl acetate. The organic

phases are combined and evaporated to dryness. The residue obtained is washed, filtered and rinsed with methanol and is then dried *in vacuo*, allowing the expected product to be isolated.

Melting point : > 350°C.

5 IR (KBr) : $\nu_{C=O} = 1691, 1749 \text{ cm}^{-1}$; $\nu_{OH, NH} = 3345 \text{ cm}^{-1}$.

Mass spectrum : 373 $[M + 1]^+$.

EXAMPLE 19 : 10-(Benzyloxy)-7-[2-(dimethylamino)ethyl][1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione

10 The expected product is obtained according to the procedure described in Example 7, starting from the compound of Example 17.

Melting point : 254°C (decomposition).

IR (KBr) : $\nu_{C=O} = 1701, 1751 \text{ cm}^{-1}$; $\nu_{NH} = 3433 \text{ cm}^{-1}$.

Mass spectrum : 520.5 $[M + 1]^+$.

15 **EXAMPLE 20 : 7-[2-(Dimethylamino)ethyl]-10-hydroxy[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione**

The expected product is obtained according to the procedure described in Example 18, starting from the compound of Example 19.

Melting point : > 360°C.

IR (KBr) : $\nu_{C=O} = 1697 \text{ cm}^{-1}$; $\nu_{OH, NH} = 3447 \text{ cm}^{-1}$.

20 Mass spectrum : 430 $[M + 1]^+$.

EXAMPLE 21 : tert-Butyl 11-(benzyloxy)-7-methyl-6,8-dioxo-7,8-dihydro[1,4]-benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-13(6H)-carboxylate

Step A : *tert-Butyl 3-[4-(1,4-benzodioxin-2-yl)-1-methyl-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl]-6-(benzyloxy)-1H-indole-1-carboxylate*

- 5 The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation L.

Melting point : 88°C.

IR (KBr) : $\nu_{\text{C=O}} = 1703, 1737 \text{ cm}^{-1}$.

Mass spectrum : 565 [M + 1]⁺.

- 10 **Step B :** *tert-Butyl 11-(benzyloxy)-7-methyl-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-13(6H)-carboxylate*

The expected product is obtained according to the procedure described in Step B of Example 17, starting from the compound obtained in Step A above.

Melting point : 204°C.

- 15 IR (KBr) : $\nu_{\text{C=O}} = 1703, 1747 \text{ cm}^{-1}$.

Mass spectrum : 463 [M-Boc + 1]⁺.

EXAMPLE 22 : 11-Hydroxy-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-carbazole-6,8(7H,13H)-dione

- 20 The expected product is obtained according to the procedure described in Step B of Example 18, starting from the compound of Example 21.

Melting point : 360°C.

IR (KBr) : $\nu_{\text{C=O}} = 1684, 1696, 1737 \text{ cm}^{-1}$; $\nu_{\text{OH,NH}} = 3317, 3442 \text{ cm}^{-1}$.

Mass spectrum : 373 [M + 1]⁺.

EXAMPLE 23 : 11-(Benzyloxy)-7-[2-(dimethylamino)ethyl][1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione

The expected product is obtained according to the procedure described in Example 7, starting from the compound of Example 21.

5 Melting point : 262°C (decomposition).

IR (KBr) : $\nu_{C=O}$ = 1701, 1750 cm^{-1} ; $\nu_{OH,NH}$ = 3432 cm^{-1} .

Mass spectrum : 520.5 $[M + 1]^+$.

EXAMPLE 24 : 7-[2-(Dimethylamino)ethyl]-11-hydroxy[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione

10 The expected product is obtained according to the procedure described in Example 18, starting from the compound of Example 23.

Melting point : > 360°C.

IR (KBr) : $\nu_{C=O}$ = 1686 cm^{-1} ; $\nu_{OH,NH}$ = 3405 cm^{-1} .

Mass spectrum : 430 $[M + 1]^+$.

15 **EXAMPLE 25 : tert-Butyl 10-(benzyloxy)-6,8-dioxo-7,8-dihydro[1,4]benzo-dioxino[2,3-a]pyrrolo[3,4-c]carbazole-13(6H)-carboxylate**

Step A : tert-Butyl 3-[4-(1,4-benzodioxin-2-yl)-1-(tert-butoxycarbonyl)-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl]-5-(benzyloxy)-1H-indole-1-carboxylate

20 The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation M.

Melting point : 76°C.

IR (KBr) : $\nu_{C=O}$ = 1741, 1763, 1797 cm^{-1} .

Mass spectrum : 551.5 [M-Boc + 1]⁺.

Step B : *tert-Butyl 10-(benzyloxy)-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-13(6H)-carboxylate*

The expected product is obtained according to the procedure described in Step B of Example 6, starting from the compound obtained in Step A above.

Melting point : > 304°C.

IR (KBr) : $\nu_{C=O}$ = 1719, 1731 cm⁻¹ ; ν_{NH} = 3198 cm⁻¹.

Mass spectrum : 549.5 [M + 1]⁺.

EXAMPLE 26 : *10-Hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-(7H,13H)-dione*

The expected product is obtained according to the procedure described in Example 8, starting from the compound of Example 25.

Melting point : > 360°C.

IR (KBr) : $\nu_{C=O}$ = 1712, 1750 cm⁻¹ ; $\nu_{NH,OH}$ = 3061, 3218, 3442 cm⁻¹.

Mass spectrum : 359 [M + 1]⁺.

EXAMPLE 27 : *tert-Butyl 11-(benzyloxy)-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-13(6H)-carboxylate*

Step A : *tert-Butyl 3-[4-(1,4-benzodioxin-2-yl)-1-(tert-butoxycarbonyl)-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl]-6-(benzyloxy)-1H-indole-1-carboxylate*

The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation N.

Melting point : 81°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1716, 1739, 1762 cm^{-1} .

Mass spectrum : 551.5 $[\text{M-Boc} + 1]^+$.

Step B : *tert-Butyl 11-(benzyloxy)-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-a]-pyrrolo[3,4-c]carbazole-13(6H)-carboxylate*

- 5 The expected product is obtained according to the procedure described in Step B of Example 6, starting from the compound obtained in Step A above.

Melting point : 199°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1731, 1761 cm^{-1} ; ν_{NH} = 3384 cm^{-1} .

Mass spectrum : 549.5 $[\text{M} + 1]^+$.

10 **EXAMPLE 28** : 11-Hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8(7H,13H)-dione

The expected product is obtained according to the procedure described in Example 8, starting from the compound of Example 27.

Melting point : > 340°C.

15 IR (KBr) : $\nu_{\text{C=O}}$ = 1711, 1745 cm^{-1} ; $\nu_{\text{OH,NH}}$ = 3054, 3197, 3427 cm^{-1} .

Mass spectrum : 359 $[\text{M} + 1]^+$.

EXAMPLE 29 : *tert-Butyl 6-methyl-5,7-dioxo-6,7-dihydronaphtho[2',3':5,6]-[1,4]dioxino[2,3-a]pyrrolo[3,4-c]carbazole-16(5H)-carboxylate*

20 **Step A** : *tert-Butyl 3-(1-methyl-4-naphtho[2,3-b][1,4]dioxin-2-yl-2,5-dioxo-2,5-dihydro-1H-pyrrol-3-yl)-1H-indole-1-carboxylate*

The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation B.

Melting point : 158°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1702, 1736 cm^{-1} .

Mass spectrum : 409 [M-Boc + 1]⁺.

5 Step B : *tert*-Butyl 6-methyl-5,7-dioxo-6,7-dihydronaphtho[2',3':5,6][1,4]dioxino-[2,3-a]pyrrolo[3,4-c]carbazole-16(5H)-carboxylate

The expected product is obtained according to the procedure described in Step B of Example 5, starting from the compound obtained in Step A above.

EXAMPLE 30 : *tert*-Butyl 8-(benzyloxy)-4-(2-hydroxyphenoxy)-1,3-dioxo-2,3-dihydropyrrolo[3,4-c]carbazole-6(1H)-carboxylate

10 The expected product is obtained in the course of purification of the compound of Step B of Example 27.

Melting point : 198°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1725, 1757 cm^{-1} ; $\nu_{\text{OH,NH}}$ = 3277 cm^{-1} .

Mass spectrum : 551.5 [M + 1]⁺.

15 EXAMPLE 31 : *tert*-Butyl 9-(benzyloxy)-4-(2-hydroxyphenoxy)-2-methyl-1,3-dioxo-2,3-dihydropyrrolo[3,4-c]carbazole-6(1H)-carboxylate

In an irradiation apparatus, 177.1 μmol of the compound obtained in Step A of Example 17 are dissolved in 500 ml of toluene and then 5.91 mmol of double-sublimed iodine are introduced into the reaction mixture. Irradiation is carried out at 500 W for 40 minutes.

20 The reaction mixture is subsequently diluted with ethyl acetate and then washed with 80 ml of 20 % aqueous sodium thiosulphate solution until the organic phase is decolourised. The aqueous phase is extracted with ethyl acetate three times, and then the organic phases are combined and directly concentrated to dryness *in vacuo*. The procedure is carried out a further three times in the same manner, using the same amounts. The crude reaction
25 products of the three procedures are combined. The residue obtained is washed, filtered

and rinsed with methanol and is then evaporated. Purification by flash chromatography over silica gel (petroleum ether/ethyl acetate : 4/6) allows the expected product to be isolated.

Melting point : 170°C.

5 IR (KBr) : $\nu_{\text{C=O}} = 1700, 1727, 1759 \text{ cm}^{-1}$; $\nu_{\text{OH}} = 3425 \text{ cm}^{-1}$.

Mass spectrum : 565.0 $[\text{M} + 1]^+$.

EXAMPLE 32 : 9-Hydroxy-4-(2-hydroxyphenoxy)-2-methylpyrrolo[3,4-c]-
carbazole-1,3(2*H*,6*H*)-dione

10 The expected product is obtained according to the procedure of Example 18, starting from the compound of Example 31.

Melting point : 277°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}} = 1698, 1757 \text{ cm}^{-1}$; $\nu_{\text{OH,NH}} = 3366 \text{ cm}^{-1}$.

Mass spectrum : 375 $[\text{M} + 1]^+$.

15 **EXAMPLE 33 :** tert-Butyl 4-(2-hydroxyphenoxy)-2-methyl-1,3-dioxo-2,3-dihydropyrrolo[3,4-c]carbazole-6(1*H*)-carboxylate

20 To a solution of 740.2 μmol of the compound of Preparation A in 12.5 ml of 1,4-dioxane there are added 616.9 μmol of the compound of Preparation E, 61.7 μmol of copper iodide and 61.7 μmol of tetrakis(triphenylphosphine)palladium. The reaction mixture is heated at 100°C, under an inert atmosphere, for 21 hours. The solution is then filtered to remove the remnants of palladium and the filtrate is evaporated to dryness. Purification by flash chromatography over silica gel (petroleum ether/ethyl acetate : 7/3) allows the expected product to be obtained.

Melting point : 182°C (decomposition).

IR (KBr) : $\nu_{\text{C=O}} = 1697, 1730, 1757 \text{ cm}^{-1}$; $\nu_{\text{OH}} = 3432 \text{ cm}^{-1}$.

25 Mass spectrum : 459 $[\text{M} + 1]^+$.

EXAMPLE 34 : **tert-Butyl 4-(2-methoxyphenoxy)-2-methyl-1,3-dioxo-2,3-dihydropyrrolo[3,4-c]carbazole-6(1H)-carboxylate**

To a solution of 152.68 μmol of the compound of Example 33 in 5 ml of acetone there are added 458.05 μmol of dry potassium carbonate and 763.41 μmol of methyl iodide. The reaction mixture is heated at reflux and stirred for 2 hours 25 minutes. The solvent is then removed *in vacuo*, and the collected residue is washed, filtered and rinsed with methanol before being dried *in vacuo*, allowing the expected product to be isolated.

Melting point : > 333°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1705, 1724 cm^{-1} .

Mass spectrum : 473.5 $[\text{M} + 1]^+$.

EXAMPLE 35 : **4-(2-Methoxyphenoxy)-2-methylpyrrolo[3,4-c]carbazole-1,3(2H,6H)-dione**

84.6 μmol of the compound of Example 34 are dissolved in 2 ml of trifluoroacetic acid and then the reaction mixture is stirred at ambient temperature for 3 hours. The acid is removed *in vacuo*, and the residue is then washed, filtered and rinsed with methanol before being dried *in vacuo*, allowing the expected product to be isolated.

Melting point : 228°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1698, 1753 cm^{-1} ; ν_{NH} = 3341, 3643 cm^{-1} .

Mass spectrum : 373 $[\text{M} + 1]^+$.

EXAMPLE 36 : **4-(2-Hydroxyphenoxy)-2-methylpyrrolo[3,4-c]carbazole-1,3-(2H,6H)-dione**

261.7 μmol of the compound of Example 33 are dissolved in 7 ml of formic acid; the reaction mixture is then stirred at ambient temperature for 18 hours. The acid is removed *in*

vacuo and the residue is then washed, filtered and rinsed with methanol before being dried *in vacuo*, allowing the expected product to be isolated.

Melting point : 258°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1695, 1757 cm^{-1} ; $\nu_{\text{OH,NH}}$ = 3243, 3525 cm^{-1} .

5 Mass spectrum : 359 $[\text{M} + 1]^+$.

EXAMPLE 37 : **tert-Butyl 2-methyl-1,3-dioxo-4-(2-{{(trifluoromethyl)sulphonyl}-oxy}phenoxy)-2,3-dihydropyrrolo[3,4-c]carbazole-6(1*H*)-carboxylate**

124.3 μmol of the compound of Example 33 are dissolved, under an anhydrous
10 atmosphere, in 1 ml of dry dichloromethane. The temperature of the solution is lowered to 0°C and then 161.5 μmol of triethylamine and 161.5 μmol of triflic anhydride are successively added dropwise to the reaction mixture, which is stirred for 1 hour 20 minutes from 0°C to ambient temperature. After hydrolysis with distilled water, the reaction mixture is diluted with dichloromethane and then the aqueous phase is extracted three
15 times with the same solvent. The organic phases are combined, dried over magnesium sulphate, filtered and concentrated. Purification by chromatography over silica gel (petroleum ether/ethyl acetate : 7/3) allows the expected product to be isolated.

Melting point : 182°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1710, 1737, 1766 cm^{-1} .

20 Mass spectrum : 591 $[\text{M} + 1]^+$.

EXAMPLE 38 : **2-[(2-Methyl-1,3-dioxo-1,2,3,6-tetrahydropyrrolo[3,4-c]carbazol-4-yl)oxy]phenyl trifluoromethanesulphonate**

Under an inert atmosphere, 67.7 μmol of the compound of Example 37 are dissolved in 1 ml of distilled dimethylformamide in the presence of 6.8 μmol of bis(triphenylphosphine)palladium chloride, 203.1 μmol of dry lithium chloride and 81.2 μmol of 1,8-diazabicyclo[5.4.0]undec-7-ene. After stirring at 140°C for 4 hours 20 minutes, the
25

reaction mixture is cooled and the dimethylformamide is then removed. Purification by chromatography over silica gel (petroleum ether/ethyl acetate : 6/4) allows the expected product to be isolated.

Melting point : 214°C.

5 IR (KBr) : $\nu_{\text{C=O}}$ = 1700-1708, 1757 cm^{-1} ; ν_{NH} = 3430 cm^{-1} .

Mass spectrum : 491 $[\text{M} + 1]^+$.

EXAMPLE 39 : **tert-Butyl 8-(benzyloxy)-4-(2-hydroxyphenoxy)-2-methyl-1,3-dioxo-2,3-dihydropyrrolo[3,4-c]carbazole-6(1H)-carboxylate**

10 The expected product is obtained in the course of purification of the compound of Step B of Example 21.

Melting point : 191°C.

IR (KBr) : $\nu_{\text{C=O}}$ = 1701, 1734 cm^{-1} ; ν_{OH} = 3425 cm^{-1} .

Mass spectrum : 565 $[\text{M} + 1]^+$.

15 **EXAMPLE 40 :** **tert-Butyl 8-(benzyloxy)-2-methyl-1,3-dioxo-4-(2-(((trifluoromethyl)sulphonyl)oxy)phenoxy)-2,3-dihydropyrrolo[3,4-c]-carbazole-6(1H)-carboxylate**

The expected product is obtained according to the procedure of Example 37, starting from the compound of Example 39.

Melting point : 186°C.

20 IR (KBr) : $\nu_{\text{C=O}}$ = 1703, 1714, 1724 cm^{-1} .

Mass spectrum : 697 $[\text{M} + 1]^+$.

EXAMPLE 41 : 8-Hydroxy-4-(2-hydroxyphenoxy)-2-methylpyrrolo[3,4-c]-carbazole-1,3(2*H*,6*H*)-dione

The expected product is obtained according to the procedure of Example 18, starting from the compound of Example 39.

5 Melting point : 200°C.

IR (KBr) : $\nu_{\text{C=O}} = 1691 \text{ cm}^{-1}$; $\nu_{\text{OH,NH}} = 3399 \text{ cm}^{-1}$.

Mass spectrum : 375 [M + 1]⁺.

EXAMPLE 42 : *tert*-Butyl 12-(benzhydryloxy)-7-methyl-6,8-dioxo-7,8-dihydro[1,4]-benzodioxino[2,3-*a*]pyrrolo[3,4-*c*]carbazole-13(6*H*)-carboxylate

10 Step A : *tert*-Butyl 7-(benzhydryloxy)-3-[4-(1,4-benzodioxin-2-yl)-1-methyl-2,5-dioxo-2,5-dihydro-1*H*-pyrrol-3-yl]-1*H*-indole-1-carboxylate

The expected product is obtained according to the procedure described in Step A of Example 5, starting from the compound of Preparation O.

Melting point : 138°C (decomposition).

15 IR (KBr) : $\nu_{\text{C=O}} = 1707, 1753 \text{ cm}^{-1}$.

Mass spectrum : 641.5 [M + 1]⁺.

Step B : *tert*-Butyl 12-(benzhydryloxy)-7-methyl-6,8-dioxo-7,8-dihydro[1,4]benzodioxino[2,3-*a*]pyrrolo[3,4-*c*]carbazole-13(6*H*)-carboxylate

20 The expected product is obtained according to the procedure described in Step B of Example 17, starting from the compound of Step A above.

PHARMACOLOGICAL STUDY OF COMPOUNDS OF THE INVENTION

EXAMPLE 43 : *In vitro* activity

✧ L1210 murine leukaemia

L1210 murine leukaemia was used *in vitro*. The cells are cultured in RPMI 1640 complete culture medium containing 10 % foetal calf serum, 2mM glutamine, 50 units/ml of penicillin, 50 µg/ml of streptomycin and 10mM Hepes, pH = 7.4. The cells are distributed on microplates and are exposed to the cytotoxic compounds for 4 doubling periods, or 48 hours. The number of viable cells is then quantified by a colorimetric assay, the Microculture Tetrazolium Assay (J. Carmichael *et al.*, Cancer Res.; 47, 939-942 (1987)). The results are expressed as the IC₅₀, the concentration of cytotoxic agent which inhibits proliferation of the treated cells by 50 %. All the compounds of the invention exhibit good cytotoxicity with respect to this cell line.

✧ Human cell lines

The compounds of the invention were also tested on human cell lines originating from solid tumours, in accordance with the same test protocol as that described for L1210 murine leukaemia but with incubation periods of 4 days instead of 2 days. By way of illustration, the compound of Example 20 has an IC₅₀ of the order of from 10 to 200nM with respect to human lines originating from prostate carcinoma, non-small-cell lung carcinoma, colon carcinoma and epidermoid carcinoma.

EXAMPLE 44 : Action on the cell cycle

L1210 cells are incubated for 21 hours at 37°C in the presence of various concentrations of test compounds. The cells are then fixed by 70 % (v/v) ethanol, washed twice in PBS and incubated for 30 minutes at 20°C in PBS that contains 100 µg/ml of RNase and 50 µg/ml of propidium iodide. The results are expressed in terms of the percentage of the cells that


accumulate in the G2+M phase after 21 hours, compared with the control (control: 20 %). The compounds of the invention are especially interesting, at a concentration of less than 2.5 μ M causing accumulation of at least 70 % of cells in the G2+M phase after 21 hours.

EXAMPLE 45 : Pharmaceutical composition : injectable solution

5	Compound of Example 20	10 mg
	Distilled water for injectable preparations	25 ml



CLAIMS

1. Compound of formula (I) :



wherein :

- 5 • A, together with the carbon atoms to which it is bonded, represents a group of formula (a) or (b) :



wherein :

- ✧ W₁, together with the carbon atoms to which it is bonded, represents a phenyl group or a pyridyl group,

5 ♣ Z represents a group selected from hydrogen and halogen atoms and the groups linear or branched (C₁-C₆)alkyl, nitro, cyano, hydroxy, linear or branched (C₁-C₆)alkoxy, aryl, aryl-(C₁-C₆)alkyl (in which the alkyl moiety is linear or branched), aryloxy and aryl-(C₁-C₆)alkoxy (in which the alkoxy moiety is linear or branched) and NR₅R₆ wherein R₅ and R₆, which are identical or different, each independently of the other represents a group selected from a hydrogen atom and a linear or branched (C₁-C₆)alkyl group,

10 ♠ R₄ represents a group selected from a hydrogen atom and the groups linear or branched (C₁-C₆)alkyl, aryl and aryl-(C₁-C₆)alkyl (in which the alkyl moiety is linear or branched) or a group -C(O)-OR'₅ wherein R'₅ represents a group selected from the groups linear or branched (C₁-C₆)alkyl, aryl and aryl-(C₁-C₆)alkyl (in which the alkyl moiety is linear or branched),

• Y represents a group selected from an oxygen atom and a methylene group,

• R₂ represents a hydrogen atom and, in that case:

15 R₃ represents a group selected from a hydrogen atom and the groups linear or branched (C₁-C₆)alkyl, aryl, aryl-(C₁-C₆)alkyl (in which the alkyl moiety is linear or branched) and SO₂CF₃,

• or R₂ and R₃ form a bond,

20 • R₁ represents a group selected from a hydrogen atom and the groups linear or branched (C₁-C₆)alkyl, aryl and aryl-(C₁-C₆)alkyl (in which the alkyl moiety is linear or branched) or a linear or branched (C₁-C₆)alkylene chain substituted by one or more identical or different groups selected from -OR''₅ and -NR''₅R''₆ wherein R''₅ and R''₆ are as defined for R₅ and R₆ defined hereinbefore,

25 • Z₁ and Z₂, which are identical or different, each independently of the other represent a hydrogen atom or


Z₁ and Z₂, together with the carbon atoms carrying them, form a phenyl group,

with the proviso that, when Z represents a hydrogen atom, R₁ is other than a hydrogen atom,

their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base,


wherein "aryl" is to be understood as meaning a phenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, indenyl or indanyl group, each of those groups being optionally substituted by one or more identical or different groups selected from halogen, linear or branched (C₁-C₆)alkyl, linear or branched (C₁-C₆)trihaloalkyl, hydroxy, linear or branched (C₁-C₆)alkoxy, and amino optionally substituted by one or two linear or branched (C₁-C₆)alkyl groups.

2. Compounds of formula (I) according to claim 1, characterised in that they represent compounds of formula (IA) :




wherein R₁, R₄, W₁, Z, Z₁ and Z₂ are as defined for formula (I), their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

3. Compounds of formula (I) according to either claim 1 or claim 2, characterised in that they represent compounds of formula (IB) :




wherein R_1 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I), their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

- 5 4. Compounds of formula (I) according to either claim 1 or claim 2, characterised in that they represent compounds of formula (IC) :




wherein R_1 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I), their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

5. Compounds of formula (I) according to claim 1, characterised in that they represent compounds of formula (ID) :




wherein R_1 , R_3 , R_4 , W_1 , Z , Z_1 and Z_2 are as defined for formula (I), their enantiomers,
 5 diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

6. Compounds of formula (I) according to either claim 1 or claim 5, characterised in that they represent compounds of formula (IE) :




10 wherein R_1 , R_3 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I), their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

7. Compounds of formula (I) according to either claim 1 or claim 5, characterised in that they represent compounds of formula (IF) :




wherein R_1 , R_3 , R_4 , Z , Z_1 and Z_2 are as defined for formula (I), their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

- 5 8. Compounds of formula (I) according to either claim 1 or claim 5, characterised in that they represent compounds of formula (IG) :




wherein R_1 , R_4 , W_1 , Z , Z_1 and Z_2 are as defined for formula (I), their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

- 10 9. Compounds of formula (I) according to either claim 1 or claim 8, characterised in that they represent compounds of formula (IH) :



wherein R₁, R₄, Z, Z₁ and Z₂ are as defined for formula (I), their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.


- 5 **10.** Compounds of formula (I) according to either claim 1 or claim 8, characterised in that they represent compounds of formula (IJ) :



wherein R₁, R₄, Z, Z₁ and Z₂ are as defined for formula (I), their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

11. Compounds of formula (I) according to any one of claims 1 to 10, characterised in that Z represents a hydrogen atom, halogen atom or hydroxy group, their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

12. Compounds of formula (I) according to any one of claims 1 to 11, characterised in that A, together with the carbon atoms to which it is bonded, represents a group of formula :



their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

13. Compounds of formula (I) according to any one of claims 1 to 12, characterised in that R₃ represents a hydrogen atom or a linear or branched (C₁-C₆)alkyl group, their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

14. Compounds of formula (I) according to any one of claims 1 to 13, characterised in that R₁ represents a hydrogen atom, a linear or branched (C₁-C₆)alkyl group or a linear or branched (C₁-C₆)alkylene chain substituted by one or more identical or different groups selected from -NR₅R₆ wherein R₅ and R₆ are as defined for formula (I), their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.


15. Compounds of formula (I) according to any one of claims 1 to 14, characterised in that Z₁ and Z₂ represent hydrogen atoms, their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.

16. Compounds of formula (I) according to claim 1, which are :

- 7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 10-fluoro-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 11-fluoro-7-methyl[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 5 • 7-[2-(dimethylamino)ethyl]-10-fluoro[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-
carbazole-6,8-dione,
- 10-hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 11-hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-dione,
- 7-[2-(dimethylamino)ethyl][1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]carbazole-6,8-
10 dione,
- 7-[2-(dimethylamino)ethyl]-10-hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-
carbazole-6,8-dione,
- 7-[2-(dimethylamino)ethyl]-11-hydroxy[1,4]benzodioxino[2,3-a]pyrrolo[3,4-c]-
carbazole-6,8-dione,
- 15 • 7-[2-(dimethylamino)ethyl][1,4]benzodioxino[2,3-e]pyrido[2',3':5,6][1,4]oxazino-
[3,2-g]isoindole-6,8-dione,

their enantiomers, diastereoisomers, N-oxide, and addition salts thereof with a pharmaceutically acceptable acid or base.


17. Process for the preparation of compounds of formula (I) according to claim 1,
20 characterised in that there is used as starting material a compound of formula (II) :



wherein Z_1 and Z_2 are as defined for formula (I),


which compound of formula (II) is reacted with N-bromosuccinimide in the presence of benzoyl peroxide to yield the compound of formula (III) :

71



wherein Z_1 and Z_2 are as defined hereinbefore,


which compound of formula (III) is reacted with sodium iodide to yield the compound of formula (IV) :



5

wherein Z_1 and Z_2 are as defined hereinbefore,

which compound of formula (IV) is reacted with n-butyllithium and then with trimethyltin chloride to yield the compound of formula (V) :




10

wherein Z_1 and Z_2 are as defined hereinbefore,


which compound of formula (V) is :

- either treated, in the presence of tetrakis(triphenylphosphine)palladium(0), with a compound of formula (VI) :




15


wherein P_G represents an amine-protecting group well-known in organic synthesis and W_1 and Z are as defined for formula (I), to yield the compound of formula (VII) :




5 wherein P_G , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
which compound of formula (VII) is treated with a compound of formula (VIII) :




10 wherein R_{1a} represents a hydrogen atom or a methyl group, to yield the compound of formula (IX) :




wherein P_G , R_{1a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
 which compound of formula (IX) is treated with N-bromosuccinimide and benzoyl
 peroxide to yield the compound of formula (I/a), a particular case of the compounds of
 formula (I) :



wherein P_G , R_{1a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
 which compound of formula (I/a) is optionally treated with a compound of
 formula (X) :




wherein R_{1b} has the same definition as R_1 in formula (I) but is other than a hydrogen
 atom or a methyl group, to yield the compound of formula (I/b), a particular case of the
 compounds of formula (I) :



wherein R_{1b}, Z, Z₁, Z₂ and W₁ are as defined hereinbefore,
 which compounds of formulae (I/a) and (I/b) constitute the compounds of
 formula (I/c) :


5




wherein R'₄ represents a hydrogen atom or a group P_G, and R₁, Z, Z₁, Z₂ and W₁ are as
 defined hereinbefore,

10

- or treated in the presence of bis(triphenylphosphine)palladium(II) chloride with a
 compound of formula (XI) :




wherein Boc represents a tert-butoxycarbonyl group and R_{1a} , W_1 and Z are as defined hereinbefore, to yield the compound of formula (XII) :




wherein Boc, R_{1a} , Z, Z_1 , Z_2 and W_1 are as defined hereinbefore, which compound of formula (XII) is :

- ◆ either irradiated with a UV lamp, in the presence of iodine, in a non-polar and aprotic solvent, to yield the compounds of formulae (I/d) and (I/e), particular cases of the compounds of formula (I) :



(I/d)




(I/e),

wherein Boc, R_{1a}, Z, Z₁, Z₂ and W₁ are as defined hereinbefore,

which compounds of formula (I/d) :

- 5 ✧ optionally are subjected to deprotection of the amine function according to conventional methods of organic synthesis to yield the compound of formula (I/f), a particular case of the compounds of formula (I) :



(I/f),


wherein R_{1a}, Z, Z₁, Z₂ and W₁ are as defined hereinbefore,

10

- ✧ or optionally are subjected to the action of a compound of formula (XIII) :




wherein R_{3a} , has the same definition as R_3 in formula (I) but is other than a hydrogen atom and G is as defined hereinbefore, to yield the compound of formula (I/g), a particular case of the compounds of formula (I) :




wherein Boc, R_{1a} , R_{3a} , Z, Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compounds of formulae (I/d), (I/e) and (I/g) constitute the compound of formula (I/h) :




wherein Boc, R_{1a} , R_3 , Z, Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compound of formula (I/h) optionally is subjected to the same reaction conditions as the compound of formula (I/a) to yield the compound of formula (I/i), a particular case of the compounds of formula (I) :



wherein R_{1b} , R_3 , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

♦ or subjected to the action of hydrochloric acid to yield the compound of formula (XIV) :




wherein R_{1a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compound of formula (XIV) is subjected to the action of a compound of formula (XV) :





wherein G represents a leaving group and R_{4a} has the same definition as R_4 in formula (I) but is other than a hydrogen atom, to yield the compound of formula (XVI) :




wherein R_{1a} , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compound of formula (XVI) is subjected to the same reaction conditions as the compound of formula (XII) to yield the compounds of formulae (I/j) and (I/k), particular cases of the compounds of formula (I) :




wherein R_{1a} , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,


which compound of formula (I/k) optionally is subjected to the action of a compound of formula (XIII) as defined hereinbefore to yield the compound of formula (I/l) :




wherein R_{1a} , R_{3a} , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
 which compounds of formulae (I/j), (I/k) and (I/l) constitute the compounds of
 formula (I/m) :



wherein R_{1a} , R_3 , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
 which compound of formula (I/m) optionally is subjected to the same reaction
 conditions as the compound of formula (I/h) to yield the compound of formula (I/n) :



wherein R_{1b} , R_3 , R_{4a} , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,
 which compounds of formulae (I/e), (I/h) and (I/i), (I/m) and (I/n) constitute the
 compounds of formula (I/o) :



wherein R_1 , R_3 , R_4 , Z , Z_1 , Z_2 and W_1 are as defined hereinbefore,

which compounds of formulae (I/a) to (I/o) constitute the totality of the compounds of
 formula (I), which are purified, where necessary, according to conventional
 purification techniques, which may be separated, if desired, into their different isomers
 according to a conventional separation technique, and which are converted, if desired,
 into their addition salts with a pharmaceutically acceptable acid or base.

18. Pharmaceutical compositions comprising, as active ingredient, at least one compound
 of formula (I) according to any one of claims 1 to 16, alone or in combination with one
 or more pharmaceutically acceptable, inert, non-toxic excipients or carriers.

19. Pharmaceutical compositions according to claim 18 for use as medicaments, in the
 treatment of cancers.

INPI

National
Institute for
Industrial Property

Received on 05/11/02

PATENT OF INVENTION
UTILITY CERTIFICATE
Intellectual Property Code - Book VI

cerfa

No. 11235*02

PATENTS DEPARTMENT
26bis, rue de Saint Pétersbourg
75800 Paris Cedex 08
Telephone: 01 53 04 53 04
Facsimile: 01 42 93 59 30

DECLARATION OF INVENTORSHIP**Page No. 1 / 3**

(if the applicant is not the inventor or not the only inventor)

This form is to be completed legibly in black ink DB 113 W /260899

Your references for this file (<i>optional</i>)		29681	
NATIONAL REGISTRATION NO.		0212965	
TITLE OF THE INVENTION (maximum 200 characters or spaces) New substituted [1,4]benzodioxino[2,3-e]isoindole compounds, a process for their preparation and pharmaceutical compositions containing them.			
APPLICANT(S): LES LABORATOIRES SERVIER 12, Place de La Défense 92415 COURBEVOIE Cedex FRANCE			
DESIGNATE(S) AS INVENTOR(S) : (Indicate at the top right-hand side "Page No. 1/1". If there are more than three inventors, use an identical form and number each page indicating the total number of pages).			
Surname		COUDERT	
Forenames		Gérard	
Address	Street	430, rue de St Denis	
	Postal code and town	45560	SAINT DENIS EN VAL (France)
Belonging company (<i>optional</i>)			
Surname		AYERBE	
Forenames		Nathalie	
Address	Street	38, rue des Aigrettes	
	Postal code and town	34250	PALAVAS LES FLOTS (France)
Belonging company (<i>optional</i>)			
Surname		LEPIFRE	
Forenames		Franck	
Address	Street	272, route d'Ardon	
	Postal code and town	45160	OLIVET (France)
Belonging company (<i>optional</i>)			
DATE AND SIGNATURE(S) OF THE APPLICANT(S) OR OF THE AUTHORISED AGENT (Name and position of signatory) 18 October 2002 [signature] Sabine WENGER, Patent Engineer			

Law No. 78-17 of 6 January 1978 relating to information processing, data files and rights applies to the responses made on this form. It guarantees right of access to and correction of the data concerning you at the INPI.

INPINational
Institute for
Industrial Property

Received on 05/11/02

**PATENT OF INVENTION
UTILITY CERTIFICATE**
Intellectual Property Code - Book VIcerfa
No. 11235*02PATENTS DEPARTMENT
26bis, rue de Saint Pétersbourg
75800 Paris Cedex 08
Telephone: 01 53 04 53 04
Facsimile: 01 42 93 59 30**DECLARATION OF INVENTORSHIP****Page No. 2 / 3**

(if the applicant is not the inventor or not the only inventor)

This form is to be completed legibly in black ink DB 113 W /260899

Your references for this file (optional)		29681	
NATIONAL REGISTRATION NO.		0212965	
TITLE OF THE INVENTION (maximum 200 characters or spaces) New substituted [1,4]benzodioxino[2,3-e]isoindole compounds, a process for their preparation and pharmaceutical compositions containing them.			
APPLICANT(S): LES LABORATOIRES SERVIER 12, Place de La Défense 92415 COURBEVOIE Cedex FRANCE			
DESIGNATE(S) AS INVENTOR(S) : (Indicate at the top right-hand side "Page No. 1/1". If there are more than three inventors, use an identical form and number each page indicating the total number of pages).			
Surname		ROUTIER	
Forenames		Sylvain	
Address	Street	140, rue d'Orléans	
	Postal code and town	45510	TIGY (France)
Belonging company (optional)			
Surname		CAIGNARD	
Forenames		Daniel-Henri	
Address	Street	22, avenue de la République	
	Postal code and town	78230	LE PECQ (France)
Belonging company (optional)			
Surname		RENARD	
Forenames		Pierre	
Address	Street	3, avenue du Parc	
	Postal code and town	78150	LE CHESNAY (France)
Belonging company (optional)			
DATE AND SIGNATURE(S) OF THE APPLICANT(S) OR OF THE AUTHORISED AGENT (Name and position of signatory) 18 October 2002 [signature] Sabine WENGER, Patent Engineer			

Law No. 78-17 of 6 January 1978 relating to information processing, data files and rights applies to the responses made on this form. It guarantees right of access to and correction of the data concerning you at the INPI.

INPI

National
Institute for
Industrial Property

Received on 05/11/02

PATENT OF INVENTION
UTILITY CERTIFICATE
Intellectual Property Code - Book VI

cerfa
No. 11235*02

PATENTS DEPARTMENT
26bis, rue de Saint Pétersbourg
75800 Paris Cedex 08
Telephone: 01 53 04 53 04
Facsimile: 01 42 93 59 30

DECLARATION OF INVENTORSHIP**Page No. 3 / 3**

(if the applicant is not the inventor or not the only inventor)

This form is to be completed legibly in black ink DB 113 W /260899

Your references for this file (<i>optional</i>)		29681	
NATIONAL REGISTRATION NO.		0212965	
TITLE OF THE INVENTION (maximum 200 characters or spaces) New substituted [1,4]benzodioxino[2,3-e]isoindole compounds, a process for their preparation and pharmaceutical compositions containing them.			
APPLICANT(S): LES LABORATOIRES SERVIER 12, Place de La Défense 92415 COURBEVOIE Cedex FRANCE			
DESIGNATE(S) AS INVENTOR(S) : (Indicate at the top right-hand side "Page No. 1/1". If there are more than three inventors, use an identical form and number each page indicating the total number of pages).			
Surname		HICKMAN	
Forenames		John	
Address	Street	136, rue de Tocqueville	
	Postal code and town	75017	PARIS (France)
Belonging company (<i>optional</i>)			
Surname		PIERRE	
Forenames		Alain	
Address	Street	9, chemin des Bois Janeaudes	
	Postal code and town	78580	LES ALLUETS LE ROI (France)
Belonging company (<i>optional</i>)			
Surname		LEONCE	
Forenames		Stéphane	
Address	Street	28B, rue Henri Simon	
	Postal code and town	78000	VERSAILLES (France)
Belonging company (<i>optional</i>)			
DATE AND SIGNATURE(S) OF THE APPLICANT(S) OR OF THE AUTHORISED AGENT (Name and position of signatory) 18 October 2002 [signature] Sabine WENGER, Patent Engineer			

Law No. 78-17 of 6 January 1978 relating to information processing, data files and rights applies to the responses made on this form. It guarantees right of access to and correction of the data concerning you at the INPI.